

www.isssi2018.org

Programme

International Symposium on

Staphylococci and Staphylococcal Infections

23 - 26 August 2018
The Maersk Tower
Copenhagen · Denmark

ISSSI
2018



Nr.	Title	Presenter	Authors
P277	Molecular Signaling Pathways leading to Ceftaroline resistance in clinical MRSA	Adriana Rosato	Maria Celeste Varela Melanie Roch Adriana Renzoni Dorte Frees Agustina Taglialegra Warren Rose Adriana Rosato
P278	Global phylogeny of Staphylococcus aureus spa type t127	Øystein Angen	Øystein Angen Patricia Alba Antonio Battisti Ivana Cirkovic Geoffrey Coombs Daum Robert D. Michael David Alessia Franco Daniel Gregson Carl Andreas Grøntvedt Thor Bech Johannesen Hulya Kaya Jesper Larsen Monica Monaco Stephan Monecke Suvi Nykäsenoja Annalisa Pantosti Frieder Schaumburg Marc Stegger Marianne Sunde Sima Tokajian Carmen Torres Deborah Williamson Anders Rhod Larsen
P279	THE PHYLOGENETIC EVOLUTION OF PORCINE MRSA IN CHINA	Stefan Schwarz	Jiang Nansong Jun Li Andrea Fessler Yang Wang Stefan Schwarz CongMing Wu
P280	Biocide susceptibility in Staphylococcus epidermidis causing infection in pets	Isabel Couto	Sofia Santos Costa Mafalda Rosa Adriana Belas Miguel Viveiros Maria Constança Pomba Isabel Couto
P281	A novel genetic locus for niche competition in staphylococci	Mal Horsburgh	Hanaa Ghabban Josephine Moran Mal Horsburgh
P282	Development of multidrug resistance in staphylococci driven by efflux	Isabel Couto	Sofia Santos Costa Adriana Rosato Miguel Viveiros Isabel Couto
P283	Prevalence of enterotoxigenic Staphylococcus aureus in cow, goat and sheep cheese in Poland	Jolanta Rola	Weronika Korpysa-Dzirba Jolanta Rola Anna Czubkowska
P284	Eradication treatment of MRSA carriage; estimation on effect of treatment and influence of PVL	Kristian Bagge	Kristian Bagge Thomas Benfield Henrik Westh Mette Damkjaer Bartels
P285	Regulation and Specificity of L _{Sa} Resistance in Staphylococci, Conferred by ABC-F Protein Vga(A)	Vladimir Vimberg	Vladimir Vimberg Lenka Nguyen Thi Ngoc Pauline Cavanagh Gabriela Balíková Novotná
P286	Diffusion and environmental persistence of pathogenic Staphylococcus capitis isolates Inside a neonatal ICU	Marine Butin	Marine Butin Yann Dumont Patricia Martins Simões Aurane Raphard Jean-Charles Picaud Frédéric Laurent

Poster Abstracts

[P282] DEVELOPMENT OF MULTIDRUG RESISTANCE IN STAPHYLOCOCCI DRIVEN BY EFFLUX

Sofia Santos Costa¹, Adriana Rosato², Miguel Viveiros³, **Isabel Couto**⁴

¹*Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa, Lisbon, Portugal*

²*Houston Methodist Research Institute, United States*

³*Unidade de Microbiologia Médica, Instituto de Higiene E Medicina Tropical, Universidade Nova de Lisboa, Loba Health and Tropical Medicine, Ihmt/Unl, Portugal, Ihmt/Unl, Portugal, Lisboa, Portugal*

⁴*Unidade de Microbiologia Médica, Global Health and Tropical Medicine (Ghtm), Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa, Lisboa, Portugal, Lisbon, Portugal*

Aim: To study the efflux driven response of two major staphylococcal pathogens, *S. aureus* and *S. epidermidis* to the challenge by non-antibiotic drugs.

Methods: We adapted three reference strains to ethidium bromide (EtBr), a broad substrate of bacterial efflux pumps. The parental strains, *S. aureus* ATCC25923, *S. epidermidis* ATCC12228 and *S. epidermidis* RP62A were cultured in varying concentrations of EtBr, to obtain their EtBr-adapted derivatives; ATCC25923_EtBr; ATCC12228_EtBr and RP62A_EtBr. Susceptibility of parental and adapted strains to 10 antibiotics and 6 biocides was evaluated by microdilution MIC determination with or without efflux inhibitors. Efflux activity was established by fluorometric assays and the relative expression of the genes coding for the main efflux pumps (EPs) of each species quantified by RT-PCR.

Results: For each strain tested, exposure to EtBr resulted in the development of a multidrug resistance (MDR) phenotype, which included resistance to fluoroquinolones and decreased susceptibility to biocides, including cetrимide, benzalkonium chloride and tetraphenylphosphonium bromide. Efflux inhibitors such as verapamil reduced these resistance levels. The EtBr-adapted cultures showed increased efflux activity, which was accompanied by over-expression of distinct EP genes, in a temporal pattern.

Conclusion: These results show that both *S. aureus* and *S. epidermidis* have the potential to develop efflux driven MDR phenotypes when exposed to a non-antibiotic substrate of multidrug EPs, which can be mediated by distinct efflux pumps, depending on the drug and the bacterial genetic background.

These results were partially presented at Costa et al BMC Microbiology, 2015,15:232; JAC, 2018, 73:320.