

1 **Compositional features and biological activities of wild and commercial**

2 ***Moringa oleifera* leaves from Guinea-Bissau**

3
4 Ângela Fernandes^{a,*}, Ângela Liberal^a, José Pinela^a, Tiane C. Finimundy^a, Aducabe Bancessi^{b,d},
5 Ana Ćirić^c, Marina Soković^c, Luís Catarino^d, Isabel C.F.R. Ferreira^a, Lillian Barros^{a,*}

6
7 ^a *Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Campus de*
8 *Santa Apolónia, 5300-253 Bragança, Portugal.*

9 ^b *Nova School of Business and Economics, NOVA University of Lisbon, Campus de Carcavelos,*
10 *Rua da Holanda, n. 1, 2775-405 Carcavelos, Portugal.*

11 ^c *Institute for Biological Research “Siniša Stanković” – National Institute of the Republic of*
12 *Serbia, University of Belgrade, Bulevar Despota Stefana 142, 11000 Belgrade, Serbia.*

13 ^d *Centre for Ecology, Evolution and Environmental Changes (cE3c), Faculty of Sciences,*
14 *University of Lisbon, Campo Grande, 1749-016 Lisbon, Portugal.*

15
16 *Corresponding authors: Â. Fernandes (afeitor@ipb.pt) and L. Barros (lillian@ipb.pt).

17 ABSTRACT

18 *Moringa oleifera* Lam. is a nutraceutical edible plant used to fight malnutrition in developing
19 countries. This investigation describes the nutritional, chemical and bioactive assets of wild and
20 commercially acquired *M. oleifera* leaves from Guinea-Bissau. Both samples presented significant
21 differences in the contents of almost all the analysed parameters. Carbohydrates and proteins were
22 the major constituents of the leaves, which also showed an interesting profile of tocopherols,
23 organic acids, unsaturated fatty acids, and phenolic compounds. The infused extract of the
24 commercial leaf sample was more effective than other preparations against lipid peroxidation and
25 oxidative haemolysis. In turn, higher antibacterial activity was achieved with hydroethanolic and
26 infused extracts of the wild sample, which extracts also displayed superior antifungal activity.
27 Overall, the commercially acquired *M. oleifera* leaves stand out with better quality profiles than
28 the wild collected, which aroused interest in studying the processing methods used locally to
29 process this functional food.

30
31 **Keywords:** *Moringa oleifera* leaves, nutritional composition, quality parameters, phenolic
32 compounds, functional foods, antioxidant/antimicrobial.

33

34 **1. Introduction**

35 In addition to their many medicinal properties, wild edible plants have been used worldwide as
36 a significant element of human nutrition (Pinela et al., 2017). Recently, plant-based functional
37 foods and nutraceuticals, as well as their potential contribution to human health and well-being,
38 have stood out in the field of scientific research (Iwu, 2017; Muyonga et al., 2020). Mostly in
39 developing countries, the food security is threatened given the exponential population growth,
40 climate changes, natural disasters, and different types of conflicts that undermine the sustainability
41 of agricultural and food systems (FAO, 2017). In West Africa, where malnutrition problems
42 persist, these trends represent a series of challenges to food and agriculture (Leakey, 2017). Thus,
43 the use of plant-based foods with nutraceutical properties, obtained from wild or easily cultivated
44 species, presents itself as a good approach to face these challenges (Gul et al., 2016).

45 *Moringa oleifera* Lam. (Fam. Moringaceae) is a fast-growing tree, native to India and Pakistan,
46 which was introduced in the tropics and sub-tropics around the world, becoming natural in several
47 African countries due to its high drought tolerance. and ability to thrive in warm and semi-arid
48 regions (Bancesi et al., 2020). This multifunctional plant is commonly used not only as a
49 vegetable, but also as a traditional remedy, as it contains a valuable nutritive profile, with
50 considerable amounts of proteins, carbohydrates, minerals, and vitamins, as well as several
51 medicinal properties. In fact, almost all portions of this plant is suitable to be exploited for food,
52 agricultural, medicinal, and industrial determinations, being an economically valued crop,
53 especially in the developing countries (Zungu et al., 2020).

54 The bark, root, leaves, and flowers of *M. oleifera* are used in several countries to prepare
55 remedies used in folk medicine to handle with skin diseases, anaemia, cholera, and further
56 illnesses. Particularly, fresh leaves are commonly consumed after cooking, or prepared in soup or

57 salads, being also employed as a food preservative to enhance the shelf-life of food, such as meat,
58 by minimizing the oxidation processes. In turn, dried leaves can be processed into powder, thus
59 being readily available for adding to foods, such as smoothies and a variety of beverages and other
60 products (Olusanya et al., 2020). *M. oleifera* leaves are also used as a good alternative to fight
61 malnutrition, especially in children, young individuals, and pregnant women, mainly due to their
62 high content of protein and dietary fibre, and their capacity to improve the quality of breastfeeding
63 milk. Besides their rich nutritional composition, *M. oleifera* leaves have also been reported to
64 contain large amounts of bioactive phytochemicals, particularly phenolic compounds, which are
65 associated with the different beneficial effects on human health (Zungu et al., 2020).

66 This deciduous tree has the potential to improve food security, human nutrition, and promote
67 rural progress and development. To maximize the benefits of domesticated plants with socio-
68 economic and technological potential for the world population, it is important to recognize how
69 they differ in compositional terms from their wild form, where the last represent a storehouse of
70 genetic diversity (Brozynska et al., 2016; Migicovsky and Myles, 2017).

71 In this sense, this study was carried out to provide a detailed characterization of the nutritional
72 profile and chemical configuration of *M. oleifera* leaves collected *in natura* and processed locally
73 (Guinea-Bissau), as well as the *in vitro* antioxidant and antimicrobial activities of their
74 hydroethanolic and aqueous extracts (prepared according to traditional practices).

75

76 **2. Material and methods**

77 *2.1. Specimen and samples preparation*

78 Wild leaf samples of *M. oleifera* (**Fig. 1a** and **b**) were collected in April 2019 in Bolama,
79 Bolama Island (11° 34'38''N 15° 28'48''W), Guinea-Bissau, and shade-dried, processed into a
80 fine powder and stowed in taped up plastic bags. Commercial leaf samples, already processed (**Fig.**
81 **1c**), were acquired in May 2019 in the Caritas Community Pharmacy in Bissau. These last samples
82 were produced and processed at Contuboel, Eastern country (12° 22'50''N 14° 33'58''W) through
83 a partnership between Caritas and a group of local women. Both samples were stored at -20 °C
84 until analysis.

85 In both sites, Bolama and Contuboel, *M. oleifera* is grown under rainfed conditions on Ferralsols
86 as living fence or near the houses in the villages. However, there are some differences between
87 these sites concerning the climate. Bolama, located at southern coastal region, receives an annual
88 rainfall of c. 2300 mm, while in Contuboel, in inner eastern country, the annual rainfall is c. 1500
89 mm. Also, the coastal influence at Bolama attenuates temperature differences and provides a less
90 dry environment in the dry season (Catarino et al., 2008)

91

92 *2.2. Nutritional value and energy determination*

93 Macronutrients amount (protein, fat, and ash; g/100 g) were assessed following the AOAC
94 official procedures of food analysis (AOAC International, 2016). The macro-Kjeldahl technique
95 was utilized to estimate the crude protein quantity ($N \times 6.25$); crude fat was determined through
96 extraction with petroleum ether in a Soxhlet apparatus; the ash totals were measured by
97 incineration at 550 ± 10 °C. The total carbohydrate sum (g/100 g of dried weight (dw)) was
98 determined by difference. The energy (kcal/100 g dw) was calculated following the equation: $4 \times$
99 (g protein + g carbohydrates) + $9 \times$ (g fat).

100 *2.3. Analysis of free sugars and organic acids*

101 The free sugars configuration was determined by high-performance liquid chromatography
102 (HPLC) coupled to a refraction index (RI) detector, by means of the internal standard (IS,
103 melezitose; Sigma-Aldrich, St. Louis, MO, USA) method, as before defined (Spréa et al., 2020).
104 The identification was performed by comparing the retention times of the authentic standards with
105 those of the samples, whereas quantification was achieved by the IS method, with calibration
106 curves built up with the standards. The results were stated in g per 100 g of dw.

107 The organic acids profile was identified by ultra-fast liquid chromatography (UFLC; Shimadzu
108 20A series, Kyoto, Japan) through a procedure previous labelled and optimized by the authors
109 (Pereira et al., 2013). Detection was done in a photo-diode array detector (PDA), using 215 nm and
110 280 nm as preferable wavelengths. Quantification was completed by likening the peak area of the
111 samples with calibration curves made with commercial standards. The organic acids quantity was
112 expressed in g per 100 g of dw.

113

114 *2.4. Analysis of fatty acids and tocopherols*

115 The fatty acid methyl esters (FAME) profile was achieved after trans-esterification of the lipid
116 fraction attained by Soxhlet extraction (Spréa et al., 2020), followed by gas-liquid chromatography
117 with flame ionization detection, using a YOUNG IN Crhomass 6500 GC System apparatus
118 equipped with a *split/splitless* injector, a flame ionization detector (FID), and a Zebron-Fame
119 column. Identification and quantification were completed by associating the relative retention
120 times of the FAME peaks of the samples, with those of the standard (47885-U; Sigma-Aldrich, St.
121 Louis, MO, USA). The Clarity DataApex 4.0 Software (Prague, Czech Republic) was utilized for
122 data handing. The results were expressed in relative percentage (%) of each detected fatty acid.

123 Tocopherols were characterized following an analytical procedure before described by the
124 authors (Spréa et al., 2020). An HPLC system coupled to a fluorescence detector (FP-2020; Jasco)
125 programed for excitation at 290 nm and emission at 330 nm was utilized. The isoforms
126 identification was attained by chromatographic comparison with authentic standards and the
127 quantification was founded on the fluorescence signal response of each standard, using the IS (tocol
128 (50 mg/mL); Matreya, Pleasant Gap, PA, USA) method and calibration curves built with
129 commercial standards. The results were expressed in mg per 100 g of dw.

130

131 *2.5. Polyphenolic profile characterization*

132 *2.5.1. Preparation of hydroethanolic and aqueous extracts*

133 The plant material was used to make hydroethanolic, infusion, and decoction extracts. The first
134 was made by stirring the plant material (~2.5 g) with 30 mL of ethanol/water (80:20, v/v) at 25 °C
135 for 1 h, and filtered through Whatman No. 4 paper. The deposit was then re-extracted with an extra
136 30 mL of the hydroalcoholic mixture. The joint extracts were concentrated at 40 °C under reduced
137 pressure (rotary evaporator Büchi R-210, Flawil, Switzerland) and further lyophilized (FreeZone
138 4.5, Labconco, Kansas City, MO, USA). For infusion extracts, 100 mL of freshly boiled distilled
139 water (heating plate, VELP scientific) were added to the plant material (~2.5 g), left aside for 5
140 min and subsequently filtered through Whatman filter paper No 4. The resultant extracts were
141 frozen and lyophilized. Decoctions were done by adding 200 mL of distilled water to the plant
142 material (~2.5 g), and boiled for 5 min. Next, the mixtures were left to rest for 5 min and then
143 filtered through Whatman No. 4 paper. The obtained decoctions were frozen and lyophilized.

144

145 *2.5.2. Analysis of phenolic compounds*

146 Phenolic compounds were investigated in the hydroethanolic, infused and decocted extracts,
147 which were redissolved in ethanol/water (80:20, v/v) and water, respectively, to a final
148 concentration of 10 mg/mL, and filtered through 0.22- μ m disposable filter disks. The analysis was
149 made in a HPLC system (Dionex Ultimate 3000 UPLC, Thermo Scientific, San Jose, CA, USA)
150 coupled with a diode-array detector (DAD, using 280 nm and 370 nm as preferred wavelengths)
151 and a Linear Ion Trap (LTQ XL) mass spectrometer (MS, Thermo Finnigan, San Jose, CA, USA)
152 prepared with an electrospray ionization (ESI) source. The separation was made in a Waters
153 Spherisorb S3 ODS-2 C18 column (3 μ m, 4.6 mm \times 150 mm; Waters, Milford, MA, USA), and
154 the elution solvents, working in the gradient, were 0.1% formic acid in water and acetonitrile. The
155 equipment and operating conditions were earlier defined by Bessada et al. (2016). Identification
156 was performed based on chromatographic behavior, spectra, and UV-Vis masses, by comparison
157 with standard compounds or data earlier described in the literature, using the Xcalibur[®] software
158 (ThermoFinnigan, San Jose, CA, USA). Quantitative analysis of the detected compounds was
159 achieved employing calibration curves based on the UV signal of the standard compounds. When
160 commercial standards were not available, the calibration curves of the most similar standards were
161 used. The phenolic standards were acquired from Extrasynthèse, Genay, France. The results were
162 expressed as mg per g of extract.

163

164 *2.6. Antioxidant activity evaluation*

165 *2.6.1. Thiobarbituric acid reactive substances (TBARS) formation inhibition capacity*

166 The lyophilized extracts were re-dissolved in water and subjected to dilutions from 2.5 mg/mL
167 to 0.0390 mg/mL. The lipid peroxidation inhibition in porcine brain cell homogenates was
168 evaluated by the reduction in TBARS; the colour strength of malondialdehyde–thiobarbituric acid

169 (MDA–TBA) was measured at 532 nm; the inhibition ratio (%) was considered using the formula:
170 $[(A - B)/A] \times 100\%$, where A and B correspond to the absorbance of the control and extract
171 sample, respectively (Spréa et al., 2020). The results were expressed in IC₅₀ values (µg/mL, sample
172 concentration providing 50% of antioxidant activity). Trolox (Sigma-Aldrich, St. Louis, MO,
173 USA) was used as positive control.

174

175 2.6.2. Oxidative haemolysis inhibition (OxHLIA) capacity

176 The antihaemolytic activity of the lyophilized extracts was evaluated by the oxidative
177 haemolysis inhibition assay (OxHLIA) using red blood cells (RBC) isolated from the blood of
178 healthy sheeps, as described by the authors (Spréa et al., 2019). The Δt values (min) resultant from
179 the half haemolysis time (Ht₅₀ values) gotten from the haemolytic curves of each extract sample
180 concentration minus the Ht₅₀ value of the PBS control, were associated to the corresponding extract
181 concentration to obtained IC₅₀ values (µg/mL), which were calculated for 60 and 120 min, *i.e.*,
182 extract concentration required to protect 50% of the erythrocyte population from the haemolytic
183 action of 2,2'-azobis(2-methylpropionamide) dihydrochloride (AAPH; Sigma-Aldrich, St. Louis,
184 MO, USA) for 60 and 120 min. Trolox was used as a positive control.

185

186 2.7. Antimicrobial activity evaluation

187 *Bacillus cereus* (food isolate), *Staphylococcus aureus* (American Type Culture Collection,
188 Manassas, VA, USA, ATCC 6538), *Listeria monocytogenes* (National Collection of Type
189 Cultures, London, UK, NCTC 7973), *Escherichia coli* (ATCC 25922), *Enterobacter cloacae*
190 (ATCC 35030), and *Salmonella typhimurium* (ATCC 13311) were selected to test the antibacterial
191 activity of the extracts. For antifungal activity, six micromycetes were used, namely *Aspergillus*

192 *fumigatus* (ATCC 9197), *Aspergillus ochraceus* (ATCC 12066), *Aspergillus niger* (ATCC 6275),
193 *Penicillium funiculosum* (ATCC 36839), *Penicillium ochrochloron* (ATCC 9112), and *Penicillium*
194 *verrucosum* var. *cyclopium* (food isolate). The microdilution method was performed as previously
195 described (Soković et al., 2010). The results were presented as the concentrations that resulted in
196 complete inhibition of the bacterial growth (MIC, minimal inhibition concentration), through the
197 colorimetric microbial viability assay, as well as MBC and MFC values (minimal bactericidal
198 concentration and minimal fungicidal concentration, respectively). The used positive controls were
199 streptomycin, ampicillin, ketoconazole, and bifonazole (Sigma-Aldrich, St. Louis, MO, USA),
200 whereas the negative control was 5% dimethyl sulfoxide (DMSO).

201

202 2.8. Statistical analysis

203 The experiments were carried out in triplicate and the results were expressed as mean \pm standard
204 deviation. the SPSS Statistics Software (IBM SPSS Statistics for Windows, Version 22.0. Armonk,
205 NY: IBM Corp.) was used to assess significant differences among the two samples by applying a
206 two-tailed paired Student's *t*-test. For phenolic compounds, differences among samples were
207 assessed applying one-way analysis of variance (ANOVA). The fulfilment of the ANOVA
208 requirements was tested through the Shapiro Wilk's and the Levene's tests to assess the normality
209 and variance homogeneity of the data, respectively. Results were compared using a Tukey's HSD
210 test. Significant differences were considered when the *p*-value was lower than 0.05.

211

212 3. Results and discussion

213 3.1. Nutritional composition of edible *M. oleifera* leaves

214 The results obtained for the *M. oleifera* leaf proximate composition are presented in **Table 1**.
215 Carbohydrates were the main constituents found in wild and commercial leaf samples (51.5 and
216 50.8 g/100 g dw, respectively), whose levels did not differ significantly ($p > 0.05$) between samples.

217 These values are slightly lower than those (56.6 g/100 g dw) reported by Ziani et al. (2019),
218 which investigate the nutritional value of *M. oleifera* cultivated in desert areas of South Algeria.
219 Contrary to the observed for carbohydrates, statistically significant differences ($p < 0.05$) were
220 found for the nutritional value constituents analysed in the wild and commercial *M. oleifera* leaf
221 samples (**Table 1**). The amount of proteins ranged from 27.4 g/100 g dw in the samples collected
222 in the wild to 32.6 g/100 g dw in the commercially acquired one. In a previous study, Chodur et
223 al. (2018) analysed domesticated and wild samples of *M. oleifera* and recorded a higher protein
224 content in the domesticated plants than in the wild collected ones (30.2 and 26.3 g/100 g,
225 respectively), although these values did not differ significantly. A higher ash content was registered
226 in the wild sample than in the commercial one (12.4 and 8.63 g/100 g dw). Valdez-Solana et al.
227 (2015) reported an appreciable amount (~11 g/100 g) of ashes in the leaves of two Mexican *M.*
228 *oleifera* cultivars, comprised within the values of this study. These results translate the high mineral
229 content that can be found in *M. oleifera* edible leaves. In turn, crude fat was the macronutrient
230 found in lower amounts, with values ranging from 8.0 to 8.7 g/100 g dw in commercial and wild
231 samples, respectively. A low amount of fat (2.82 g/100 g) was also found by Ilyas et al. (2015),
232 which demonstrates that *M. oleifera* leaves can be suitable foods for low-fat diets, with an energy
233 contribution ranging from 394 to 406 kcal/100 g dw (**Table 1**).

234 **Table 1** presents the free sugars composition of *M. oleifera* leaves. Fructose, glucose, sucrose,
235 and trehalose were detected in both samples and sucrose was the major free sugar, with levels
236 varying significantly ($p < 0.001$) from 1.97 g/100 g dw in the wild sample to 5.33 g/100 g dw in

237 the commercial one. In a previous study, Al Juhaimi et al. (2017) reported small amounts of
238 raffinose and stachyose in *M. oleifera* leaves, but did not detect trehalose. The authors quantified
239 glucose as the most abundant free sugar in their samples (6.65 g/100 g), followed by fructose (3.96
240 g/100 g), a result that differs from this study where sucrose was the most abundant free sugar. All
241 these variances might be owed to edaphoclimatic influences, such as temperature variations and
242 soil composition, as well as some biotic conditions that can influence the synthesis of sugars.

243 Concerning organic acids, oxalic, malic, ascorbic, and citric acids were identified in the studied
244 leaf samples (**Table 1**). Malic acid, a dicarboxylic acid that contributes to a sour taste and used as
245 a food additive was the most abundant (~4 g/100 g dw). The levels of each identified organic acid
246 differed significantly ($p \leq 0.05$) between samples. The observed differences may be related to the
247 maturity of the leaves at harvest, drying and storage practices and the processing conditions applied
248 to the samples (Dias et al., 2016). Previously, Ziani et al. (2019) identified three organic acids in
249 *M. oleifera* leaves, namely oxalic, malic, and ascorbic acids; malic acid was also detected in higher
250 amounts (2.5 g/100g dw), followed by oxalic and ascorbic acids. The authors did not identify citric
251 acid (a natural food additive), although it was found in considerable amounts in our samples, a
252 result that supports the use of this plant as a traditional food preservative.

253 The fatty acids profile results are presented in **Table 2**. A total of 21 compounds were identified
254 in the analysed *M. oleifera* leaves. α -Linolenic acid (C18:3n3) was the most abundant, followed
255 by palmitic (C16:0) and linoleic (C18:2n6) acids. C16:0 was the only fatty acid that did not show
256 significant differences ($p > 0.05$) between samples. Considering their classification,
257 polyunsaturated fatty acids (PUFA) represented the higher percentage of all fatty acids, with values
258 ranging from 58.5 to 58.8 %, followed by saturated fatty acids (SFA) and monounsaturated fatty
259 acids (MUFA). Both leaf samples presented significant differences in the contents of SFA and

260 MUFA. Castillo-López et al. (2017) in a study that aimed to characterize two *M. oleifera* leaf
261 variants that grow in Sinaloa, Mexico, identified 14 fatty acids, where the α -linolenic acid was
262 found in higher amounts (ranging from 62.72 to 66.79 %), followed by palmitic and linoleic acids,
263 being PUFA the detached class of fatty acids in both samples. According to various studies, the
264 consumption of PUFA is responsible for decreasing the levels of total and low-density lipoprotein
265 (LDL) cholesterol, playing a cardioprotective role in human organisms (Ander et al., 2003).

266 Regarding tocopherols, the isoforms α -, β - and δ -tocopherol were identified in both wild and
267 commercial leaf samples (**Table 2**) and α -tocopherol was the main isoform (with 58.79 and 68.03
268 mg/100 g dw, respectively), followed by δ -tocopherol, and lastly β -tocopherol. Therefore, a 100 g
269 serving of *M. oleifera* leaves provides more vitamin E than the recommended dietary allowances
270 (15 mg/day) for adult individuals. It was also observed that the origin of the plant material caused
271 significant differences ($p < 0.001$) in the content of these lipophilic antioxidants. A study of Ziani
272 et al. (2019) describes the four tocopherol isoforms in amounts different than those found in this
273 study; the authors also reported α -tocopherol as the main isoform (quantified in lower amounts of
274 4.0 mg/100 g dw), followed by γ -tocopherol (with just 0.41 mg/100 g dw). These compositional
275 differences may be due to the edaphoclimatic conditions of the growing sites, abiotic factors, and
276 the processing methods used in samples preparation.

277

278 3.2. Polyphenolic composition of *M. oleifera* leaf extracts

279 The data attained regarding the chromatographic features (retention time, λ_{\max} , pseudomolecular
280 ions, and main MS² fragments), as well as the tentative identification of the phenolic compounds
281 found in each hydroethanolic, infused and decocted extract of *M. oleifera* leaves are shown in
282 **Table 3**. Twenty-five compounds were identified, 6 of which were phenolic acids and 19 derived

283 from glycosylated flavonols. Regarding the phenolic acids and considering the above parameters,
284 the compounds were tentatively identified as caffeic acid hexoside (peak **1**; λ_{\max} , 324 nm; $[\text{M-H}]^-$
285 at m/z 341), 3-*O*- and 4-*O*-caffeoylquinic acids (peaks **2** and **3**; λ^{\max} , 324 nm; $[\text{M-H}]^-$ at m/z 353,
286 respectively), 3-*p*-coumaroylquinic acid (peak **4**; λ_{\max} , 280 nm; $[\text{M-H}]^-$ at m/z 337), and finally *cis*
287 and *trans* 5-*O*-caffeoylquinic acids (peak **5** and **6**, respectively; λ_{\max} , 324 nm; $[\text{M-H}]^-$ at m/z 353).
288 Chlorogenic acids are a large family of polyphenols that function as antioxidants and as dietary
289 neurosignalling compounds, improving lipid metabolism and promoting weight loss by reducing
290 the synthesis of visceral fat, cholesterol, and fatty acids (Nakatani et al., 2000). These phenolic
291 acids are composed of quinic acid linked to *trans*-cinnamic acids, such as caffeic acid, via an ester
292 bond.

293 Although the most common chlorogenic acid is 5-*O*-caffeoylquinic, other isomers may be
294 present in the same plant, such as 3 and 4-*O*-caffeoylquinic acids (Braham et al., 2020a). Since *M.*
295 *oleifera* has been widely studied through the last decades, the compounds found in our study have
296 previously been identified in the foliar parts of this species from Brazil, namely the caffeic acid
297 (Oldoni et al., 2019), and from South Algeria decocted and hydroethanolic extracts (peaks 2, 3 and
298 4) by Ziani et al. (2019). However, our study also reveals the presence of *cis* and *trans* 5-*O*-
299 caffeoylquinic acid compounds, while Braham et al. (2020) only used this chlorogenic acid as a
300 pattern (5-*O*-caffeoylquinic acid) to identify the respective isomers. In fact, and contrary to our
301 study, Castro-López et al. (2017) reported that the only chlorogenic acid present in *M. oleifera* leaf
302 extracts obtained by decoction is the 3-*O*-caffeoylquinic acid.

303 The group of flavonoids proved to be the most abundant in the prepared *M. oleifera* leaf extracts,
304 among which stands out the quercetin derivatives, present in greater number than the other
305 identified aglycone flavonoids, followed by apigenin, isorhamnetin and kaempferol derivatives in

306 glycosidic and isomeric forms. Thus, our experimental conditions allow us to identify quercetin-
307 3-*O*-rhamnoside (peak **11**; [M-H]⁻ at *m/z* 447), quercetin-3-*O*-rutinoside (peak **13**; [M-H]⁻ at *m/z*
308 609), quercetin 3-*O*-glucoside (peak **16**; [M-H] at *m/z* 463), and quercetin-3-*O*-(6''-malonyl-
309 glucoside) (peak **18**; [M-H] at *m/z* 549), already previously identified in *M. oleifera* leaves from
310 Southern Italy (Rocchetti et al., 2020) and Ghana (Amaglo et al., 2010a), respectively, and the last
311 two compounds by Bennett et al. (2003) from different locations of Africa. Peaks **14/15**, presented
312 both a pseudomolecular ion [M-H]⁻ at *m/z* 609, and MS² fragment at *m/z* 301, allowing its tentative
313 identification as quercetin-4-*O*-rhamnoside-7-*O*-hexoside and quercetin-3-*O*-rhamnoside-7-*O*-
314 hexoside, respectively, firstly identified in *M. oleifera* leaves in our study.

315 Given this assumption, its tentative identification was achieved following the earlier described
316 by Carocho et al. (2014) in *Castanea sativa* Mill. flower extracts. Likewise, peak **17** ([M-H]⁻ at
317 *m/z* 505) and peaks **19-21** ([M-H] at *m/z* 549) were tentatively identified as quercetin-3-*O*-(6''-
318 acetyl)-glucoside (Jang et al., 2018), quercetin 7-*O*-malonylglucoside (Guimarães et al., 2014),
319 quercetin-*O*-malonyl-hexoside (Bessada et al., 2016), and quercetin 3-*O*-malonylglucoside (Lin et
320 al., 2020), as described by the authors. A total of 6 quercetin derivatives were, to the best of our
321 knowledge, identified for the first time in *M. oleifera* leaf extracts, probably owing to specific
322 environmental characteristics and geographic location in which this species grows (Ma et al.,
323 2020). Quercetin is described in the literature for its high antioxidant capacity, given its high ability
324 to eliminate reactive oxygen species, and for its antitumor properties, since it has the ability to
325 modulate cell apoptosis and the migration and growth of tumour cells through the various
326 signalling pathways involved (Lesjak et al., 2018). Apigenin-derived compounds appear as the
327 second largest group of flavonoids herein identified, with peaks **7-10** ([M-H]⁻ at *m/z* 593)
328 tentatively identified as apigenin-6,8-di-*C*-hexoside, apigenin 8-*C*-glucoside-6''-*O*-glucoside,

329 apigenin-6-*C*-glucoside-7-*O*-glucoside, and apigenin-6,8-*C*-di- β -*D*-glucopyranoside, respectively,
330 following the before defined by Ferreres et al. (2007), and once again herein identified for the first
331 time in *M. oleifera* species. On the other hand, peak **12** ($[M-H]^-$ at m/z 431), tentatively identified
332 as apigenin-8-*C*-glucoside, was previously reported by Karthivashan et al. (2013) in extracts with
333 different hydroethanolic gradients of *M. oleifera* leaves from Malaysia.

334 Three isorhamnetin derivatives were also identified in our study, namely isorhamnetin-3-*O*-
335 rutinoside (peak **22**; $[M-H]^-$ at m/z 623), isorhamnetin-3-*O*-(6"-acetyl)-glucoside (peak **24**; $[M-H]^-$
336 at m/z 519), and isorhamnetin-3-*O*-(6''-malonyl)-glucoside (peak **25**; $[M-H]^-$ at m/z) 563, being
337 the first two identified in *M. oleifera* leaf samples from South Africa by Makita et al. (2016) and
338 the last compound (peak 25) by Amaglo et al. (2010) in samples from Ghana. The only kaempferol
339 derivate tentatively identified in our study, namely kaempferol-3-*O*-(6''-malonyl-glucoside) (peak
340 **23**; $[M-H]^-$ at m/z 533), was also identified in the same samples of *M. oleifera* leaves from Ghana
341 (Amaglo et al., 2010b).

342 **Table 4** presents the quantitative data of the phenolic compounds tentatively identified in *M.*
343 *oleifera* leaves. Each group of *M. oleifera* leaf extracts presented a dissimilar profile of the
344 identified phenolic compounds, both in quantitative and qualitative terms. Therefore, it is difficult
345 to point out the best extract preparation method to obtain a higher quantity of phenolic compounds.
346 However, five of the identified compounds were present in all the prepared extracts, namely the
347 compounds corresponding to the peaks **2**, **5**, **13**, **14**, and **18**. It is well supported by the literature
348 that the extraction method (and solvent) used to recover phenolic compounds directly influences
349 both the extraction yield and the phytochemical profile of the samples under analysis (Celep et al.,
350 2019a), which explains the quantitative differences observed in this study.

351 The hydroethanolic extract of the wild *M. oleifera* leaf sample showed the main total
352 concentration of phenolic compounds (103.7 ± 0.2 mg/g of extract), followed by the infused
353 (51.8 ± 0.5 mg/g of extract) and decocted (47 ± 1 mg/g of extract) extracts of the same plant material,
354 which can be explained by the use of different extraction solvents (ethanol and water) or by the
355 application of heat in the preparation of the aqueous extracts (Celep et al., 2019b). On the other
356 hand, the same extraction trend was not observed in the commercially acquired sample, since the
357 higher concentration of phenolic compounds (72 ± 1 mg/g of extract) was found in the decocted
358 extract, followed by hydroethanolic (65.3 ± 0.3 mg/g of extract) and infused (52 ± 1 mg/g of extract)
359 extracts. This contrast in the total content of phenolic compound in the prepared extracts, and
360 between wild collected and commercial acquired samples may be attributed to the processing and
361 preparation methods already mentioned above to which the samples were subjected (Dias et al.,
362 2016).

363 Despite the different phenolic profiles found between each of the extracts and the wild and
364 commercial *M. oleifera* leaf samples, the group of flavonoids stands out in both cases, which is in
365 agreement with the study of Ziani et al. (2019), whose investigation exposes a total concentration
366 of flavonoids of up to 30 mg/g extract.

367

368 3.3. Biological activities of *M. oleifera* leaf extracts

369 Wild and commercial *M. oleifera* leaf hydroethanolic, infused, and decocted extract were tested
370 for their capacity to prevent lipid peroxidation, using porcine brain cell tissues as oxidizable
371 substrates, and the oxidative haemolysis using sheep RBC as a model. The results of both *in vitro*
372 assays are presented in **Table 5**. Regarding the TBARS assay, the hydroethanolic extract of both
373 wild and commercial samples showed the best antioxidant activity (IC_{50} values of 54 and 57

374 $\mu\text{g/mL}$) when compared with the aqueous extracts, and with no statistical difference between
375 values. The lowest TBARS formation inhibition capacity (higher IC_{50} values) was achieved with
376 the extract prepared by decoction, especially from leaves collected in the wild. The different
377 solvents, temperatures, and extraction times may justify these differences, since certain antioxidant
378 compounds may have been better extracted or degraded by these processing variables.

379 A different trend was observed in the OxHLIA assay (**Table 5**); the aqueous extracts were more
380 effective in protecting 50% of the RBC population from the AAPH-induced oxidative haemolysis.
381 In fact, these extracts were more active than the synthetic antioxidant trolox used as a positive
382 control, which is a very promising result for a natural plant extract. While the commercial leaf
383 sample yielded lower IC_{50} values when prepared in infusion, the wild sample was more antioxidant
384 when extracted by decoction. In addition, while both aqueous extracts exerted protective effects
385 up to periods of 120 min, the hydroethanolic extract just protected the RBC for a 60 min Δt .
386 The antioxidant activity of *M. oleifera* leaf extract has been reported by some authors. Chodur et
387 al. (2018) evaluate the antioxidant activity of wild and domesticated *M. oleifera* samples and found
388 that the last ones presented higher activity than the “wild type”, which in agreement with the results
389 herein obtained for the TBARS assay. An earlier report of Ilyas *et al.* (2015) also attributes a strong
390 antioxidant activity to *M. oleifera* leaf powder when tested for DPPH radical scavenging activity.

391 The results of the antibacterial activity of the three *M. oleifera* leaf extracts are presented in
392 **Table 6**. This activity was screened against six foodborne bacteria, namely *Bacillus cereus*,
393 *Staphylococcus aureus*, *Listeria monocytogenes*, *Escherichia coli*, *Enterobacter cloacae*, and
394 *Salmonella typhimurium*, against which strong inhibitory and bactericidal effects were observed,
395 in most cases higher than the MIC and MBC values of the antibiotics streptomycin and ampicillin.
396 The decoctions of leaves collected in the wild showed higher MIC and MBC values than the other

397 two extracts (except for *E. cloacae*), thus requiring a higher extract concentration to inhibit or kill
398 the tested bacterial strains. For commercial samples, infusions were the less effective against *S.*
399 *aureus*, *E. cloacae*, *S. typhimurium*, yet the MIC/MBC values of these last two Gram-negative
400 bacteria were lower or equal to those of the positive controls. Overall, wild samples appeared to
401 have a better antibacterial capacity than the commercially acquired ones. A previous publication
402 of Bukar et al. (2010) also describes that the *M. oleifera* ethanolic leaf extract exhibits a broad-
403 spectrum activity against foodborne pathogens, such as *E. coli*, *S. aureus*, *Pseudomonas*
404 *aeruginosa*, and *Enterobacter aerogenes*.

405 **Table 6** also present the antifungal activity results achieved with the *M. oleifera* leaf extracts
406 against *Aspergillus fumigatus*, *A. ochraceus*, *A. niger*, *Penicillium funiculosum*, *P. ochrochloron*,
407 and *P. verrucosum* var. *cyclopium*. In general, the plant extracts exhibited strong inhibitory and
408 fungicidal potential when compared to the positive controls ketoconazole and bifonazole. As
409 observed for antibacterial activity, infusions from the commercial sample were the least effective
410 preparations against all tested fungi.

411

412 **4. Conclusion**

413 The present investigation emphasised the nutritional quality and bioactive potential of wild
414 harvested and commercially acquired *M. oleifera* leaves. This functional food is rich in
415 carbohydrates (including sucrose, fructose, and trehalose), proteins, minerals, and PUFA (mainly
416 C18:3n3 and C18:2n6), and vitamin E, due to the high contents of α -tocopherol. It also contains
417 high levels of malic and citric acids, which may import certain organoleptic features, as well as
418 ascorbic acid. The HPLC-DAD-ESI/MSⁿ analysis allowed identifying flavonoids (comprising
419 apigenin, quercetin, isorhamnetin, and kaempferol glycoside derivatives) as the major class of

420 polyphenols in hydroethanolic, infused, and decocted leaf extracts, followed by phenolic acids
421 such as caffeoylquinic acids. The hydroethanolic and aqueous preparations were capable of
422 inhibiting the TBARS formation and the oxidative haemolysis, and were effective against some
423 food-borne bacterial and fungal strains.

424 The commercially acquired *M. oleifera* leaves show up with better quality composition than the
425 wild collected ones, highlighting the importance of processing methods in the nutritional and
426 chemical profile of this plant.

427 Altogether, these results highpoint the nutritional value of this low-cost functional food widely
428 available in Guinea-Bissau and other tropical regions around the world, their suitability to fight
429 malnutrition and increase food security in developing countries, and potential to be used in the
430 development of nutraceutical formulations.

431

432 **Acknowledgements**

433 The authors are thankful to the Foundation for Science and Technology (FCT, Portugal) for
434 financial support through national funds FCT/MCTES to CIMO (UIDB/00690/2020) and to the
435 Centre for Ecology, Evolution and Environmental Changes (cE3c) (UIDB/00329/2020). National
436 funding by FCT, P.I., through the institutional scientific employment program-contract for the
437 contracts of A. Fernandes and L. Barros, and through the individual scientific employment
438 program-contract for the contract of J. Pinela (CEECIND/01011/2018). To the grants of Â. Liberal
439 (through the INTERNOVAMARKET-FOOD project) and A. Bancessi (SFRH/BD/135356/2017).

440 This work was funded by FEDER-Interreg España-Portugal programme through the projects
441 0377_Iberphenol_6_E and TRANSCoLAB 0612_TRANS_CO_LAB_2_P, and also by the
442 Ministry of Education, Science and Technological Development of Republic of Serbia (451-03-
443 68/2020-14/200007).

444

445 **CRedit authorship contribution statement**

446 **Ângela Fernandes:** Formal analysis, Methodology, Software, Validation, Investigation, Data
447 curation, Writing - original draft. **Ângela Liberal:** Writing - original draft. **José Pinela:**
448 Methodology, Software, Validation, Investigation, Methodology, Data curation, Writing - review
449 & editing. **Tiane C. Finimundy:** Methodology, Software, Validation, Investigation, Data curation.
450 **Aducabe Bancesi:** Investigation, Writing - review & editing. **Ana Ćirić:** Investigation, Formal
451 analysis. **Marina Soković:** Investigation, Formal analysis; **Luís Catarino:** Supervision, Writing -
452 review & editing. **Isabel C.F.R. Ferreira:** Supervision, Project administration. **Lillian Barros:**
453 Conceptualization, Validation, Investigation, Writing - review & editing, Visualization,
454 Supervision, Project administration, Funding acquisition.

455

456 **Declarations of interest**

457 The authors have declared no conflict of interest.

458

459

460 **References**

461 Al Juhaimi, F., Ghafoor, K., Babiker, E.E., Matthäus, B., Özcan, M.M., 2017. The biochemical
462 composition of the leaves and seeds meals of *Moringa* species as non-conventional sources
463 of nutrients. J Food Biochem. 41, e12322.

464 Amaglo, N.K., Bennett, R.N., Lo Curto, R.B., Rosa, E.A.S., Lo Turco, V., Giuffrida, A., Curto, A.
465 Lo, Crea, F., Timpo, G.M., 2010a. Profiling selected phytochemicals and nutrients in different
466 tissues of the multipurpose tree *Moringa oleifera* L., grown in Ghana. Food Chem. 122, 1047–
467 1054.

468 Amaglo, N.K., Bennett, R.N., Lo Curto, R.B., Rosa, E.A.S., Lo Turco, V., Giuffrida, A., Curto, A.
469 Lo, Crea, F., Timpo, G.M., 2010b. Profiling selected phytochemicals and nutrients in different
470 tissues of the multipurpose tree *Moringa oleifera* L., grown in Ghana. Food Chem. 122, 1047–
471 1054.

472 Ander, B.P., Dupasquier, C.M.C., Prociuk, M.A., Pierce, G.N., 2003. Polyunsaturated fatty acids
473 and their effects on cardiovascular disease. Exp Clin Cardiol. 8, 164-172.

474 AOAC International, 2016. Official Methods of Analysis of AOAC International, 20th ed. AOAC
475 International, Gaithersburg.

476 Bancessi, A., Bancessi, Q., Baldé, A., Catarino, L., 2020. Present and potential uses of *Moringa*
477 *oleifera* as a multipurpose plant in Guinea-Bissau. S. Afr. J. Bot. 129, 206–208.

478 Bennett, R.N., Mellon, F.A., Foidl, N., Pratt, J.H., Dupont, M.S., Perkins, L., Kroon, P.A., 2003.
479 Profiling glucosinolates and phenolics in vegetative and reproductive tissues of the multi-
480 purpose trees *Moringa oleifera* L. (Horseradish tree) and *Moringa stenopetala* L. Journal of
481 Agricultural and Food Chem. 51, 3546–3553.

482 Bessada, S.M.F., Barreira, J.C.M., Barros, L., Ferreira, I.C.F.R., Oliveira, M.B.P.P., 2016.
483 Phenolic profile and antioxidant activity of *Coleostephus myconis* (L.) Rchb.f.: An
484 underexploited and highly disseminated species. Ind Crops Prod.89, 45–51.

485 Braham, F., Carvalho, D.O., Almeida, C.M.R., Zaidi, F., Magalhães, J.M.C.S., Guido, L.F.,
486 Gonçalves, M.P., 2020a. Online HPLC-DPPH screening method for evaluation of radical
487 scavenging phenols extracted from *Moringa oleifera* leaves. S. Afr. J. Bot. 129, 146–154.

488 Braham, F., Carvalho, D.O., Almeida, C.M.R., Zaidi, F., Magalhães, J.M.C.S., Guido, L.F.,
489 Gonçalves, M.P., 2020b. Online HPLC-DPPH screening method for evaluation of radical
490 scavenging phenols extracted from *Moringa oleifera* leaves. S. Afr. J. Bot. 129, 146–154.

491 Brozynska, M., Furtado, A., Henry, R.J., 2016. Genomics of crop wild relatives: Expanding the
492 gene pool for crop improvement. *Plant Biotechnol J.* 14, 1070–1085.

493 Bukar, A., Uba, A., Oyeyi, T., 2010. Antimicrobial profile of *Moringa oleifera* lam. extracts
494 against some food-borne microorganisms. *Bayero J Pure Applied Sci.* 3, 43–48.

495 Carocho, M., Barros, L., Bento, A., Santos-Buelga, C., Morales, P., Ferreira, I.C.F.R., 2014.
496 *Castanea sativa* Mill. Flowers amongst the most powerful antioxidant matrices: a
497 phytochemical approach in decoctions and infusions. *Biomed Res Int.* 2014, 232956.

498 Castillo-López, R.I., León-Felix, J., Angulo-Escalante, M., Gutiérrez-Dorado, R., Muy-Rangel,
499 M.D., Heredia, J.B., 2017. Nutritional and phenolic characterization of *Moringa oleifera*
500 leaves grown in Sinaloa, México. *Pak J Bot* 49, 161–168.

501 Castro-López, C., Ventura-Sobrevilla, J.M., González-Hernández, M.D., Rojas, R., Ascacio-
502 Valdés, J.A., Aguilar, C.N., Martínez-Ávila, G.C.G., 2017. Impact of extraction techniques
503 on antioxidant capacities and phytochemical composition of polyphenol-rich extracts. *Food*
504 *Chem.* 237, 1139–1148.

505 Celep, E., Seven, M., Akyüz, S., İnan, Y., Yesilada, E., 2019a. Influence of extraction method on
506 enzyme inhibition, phenolic profile and antioxidant capacity of *Sideritis trojana* Bornm. *S.*
507 *Afr. J. Bot.* 121, 360–365.

508 Celep, E., Seven, M., Akyüz, S., İnan, Y., Yesilada, E., 2019b. Influence of extraction method on
509 enzyme inhibition, phenolic profile and antioxidant capacity of *Sideritis trojana* Bornm. *S.*
510 *Afr. J. Bot.* 121, 360–365.

511 Chodur, G.M., Olson, M.E., Wade, K.L., Stephenson, K.K., Nouman, W., Garima, Fahey, J.W.,
512 2018. Wild and domesticated *Moringa oleifera* differ in taste, glucosinolate composition, and
513 antioxidant potential, but not myrosinase activity or protein content. *Sci Rep.* 8, 7995.

514 Dias, M.I., Barros, L., Morales, P., Cámara, M., Alves, M.J., Oliveira, M.B.P.P., Santos-Buelga,
515 C., Ferreira, I.C.F.R., 2016. Wild *Fragaria vesca* L. fruits: a rich source of bioactive
516 phytochemicals. *Food Funct.* 7, 4523–4532.

517 FAO, 2017. The future of food and agriculture: Trends and challenges.

518 Ferreres, F., Gil-Izquierdo, A., Andrade, P.B., Valentão, P., Tomás-Barberán, F.A., 2007.
519 Characterization of C-glycosyl flavones O-glycosylated by liquid chromatography-tandem
520 mass spectrometry. *J. Chromatogr. A* 1161, 214–223.

521 Guimarães, R., Barros, L., Calhella, R.C., Carvalho, A.M., Queiroz, M.J.R.P., Ferreira, I.C.F.R.,
522 2014. Bioactivity of different enriched phenolic extracts of wild fruits from northeastern
523 Portugal: a comparative study. *Plant Foods Hum Nutr.* 69, 37-42.

524 Gul, K., Singh, A.K., Jabeen, R., 2016. Nutraceuticals and functional foods: the foods for the future
525 world. *Crit Rev Food Sci.* 56, 2617–2627.

526 Ilyas, M., Arshad, M.U., Saeed, F., Iqbal, M., 2015. Antioxidant potential and nutritional
527 comparison of moringa leaf and seed powders and their tea infusions. *J Anim Plant Sci.* 25,
528 226–233.

529 Iwu, M.M., 2017. *Food as Medicine: Functional Food Plants of Africa - 1st Edition - Mau.* CRC
530 Press.

531 Jang, G.H., Kim, H.W., Lee, M.K., Jeong, S.Y., Bak, A.R., Lee, D.J., Kim, J.B., 2018.
532 Characterization and quantification of flavonoid glycosides in the *Prunus* genus by UPLC-
533 DAD-QTOF/MS. *Saudi J Biol Sci.* 25, 1622–1631.

534 Karthivashan, G., Tangestani Fard, M., Arulselvan, P., Abas, F., Fakurazi, S., 2013. Identification
535 of bioactive candidate compounds responsible for oxidative challenge from hydro-ethanolic
536 extract of *Moringa oleifera* leaves. *Int. J. Food Sci.* 78, C1368-1375.

537 Leakey, R., 2017. Multifunctional Agriculture: Achieving Sustainable Development in Africa.
538 Academic Press.

539 Lesjak, M., Beara, I., Simin, N., Pintać, D., Majkić, T., Bekvalac, K., Orčić, D., Mimica-Dukić,
540 N., 2018. Antioxidant and anti-inflammatory activities of quercetin and its derivatives. J.
541 Funct. Foods. 40, 68–75.

542 Lin, Y.C., Wu, C.J., Kuo, P.C., Chen, W.Y., Tzen, J.T.C., 2020. Quercetin 3-O-malonylglucoside
543 in the leaves of mulberry (*Morus alba*) is a functional analog of ghrelin. J Food Biochem. 44,
544 e13379.

545 Ma, Z.F., Ahmad, J., Zhang, H., Khan, I., Muhammad, S., 2020. Evaluation of phytochemical and
546 medicinal properties of Moringa (*Moringa oleifera*) as a potential functional food. S. Afr. J.
547 Bot. 129, 40–46.

548 Makita, C., Chimuka, L., Steenkamp, P., Cukrowska, E., Madala, E., 2016. Comparative analyses
549 of flavonoid content in *Moringa oleifera* and *Moringa ovalifolia* with the aid of UHPLC-
550 qTOF-MS fingerprinting. S. Afr. J. Bot. 105, 116–122.

551 Migicovsky, Z., Myles, S., 2017. Exploiting wild relatives for genomics-assisted breeding of
552 perennial crops. Front. Plant Sci. 8, 460.

553 Muyonga, J.H., Nansereko, S., Steenkamp, I., Manley, M., Okoth, J.K., 2020. Traditional African
554 foods and their potential to contribute to health and nutrition: traditional African foods, in:
555 African Studies: Breakthroughs in Research and Practice. p. 27.

556 Nakatani, N., Kayano, S.I., Kikuzaki, H., Sumino, K., Katagiri, K., Mitani, T., 2000. Identification,
557 quantitative determination, and antioxidative activities of chlorogenic acid isomers in prune
558 (*Prunus domestica* L.). J. Agric. Food Chem. 48, 5512–5516.

559 Oldoni, T., Merlin, N., Karling, M., Carpes, S., Alencar, C., Morales, R., da Silva, E., Pilau, E.,
560 2019. Bioguided extraction of phenolic compounds and UHPLC-ESI-Q-TOF-MS/MS
561 characterization of extracts of *Moringa oleifera* leaves collected in Brazil. *Food Res. Int.* 125,
562 108647.

563 Olusanya, R.N., Kolanisi, U., van Onselen, A., Ngobese, N.Z., Siwela, M., 2020. Nutritional
564 composition and consumer acceptability of *Moringa oleifera* leaf powder (MOLP)-
565 supplemented *mahewu*. *S. Afr. J. Bot.* 129, 175–180.

566 Pereira, C., Barros, L., Carvalho, A.M., Ferreira, I.C.F.R., 2013. Use of UFLC-PDA for the
567 analysis of organic acids in thirty-five species of food and medicinal plants. *Food Anal*
568 *Methods.* 6, 1337–1344.

569 Pinela, J., Carvalho, A.M., Ferreira, I.C.F.R., 2017. Wild edible plants: Nutritional and
570 toxicological characteristics, retrieval strategies and importance for today's society. *Food*
571 *Chem. Toxicol.* 110, 165–188.

572 Rocchetti, G., Pagnossa, J.P., Blasi, F., Cossignani, L., Hilsdorf Piccoli, R., Zengin, G.,
573 Montesano, D., Cocconcelli, P.S., Lucini, L., 2020. Phenolic profiling and in vitro bioactivity
574 of *Moringa oleifera* leaves as affected by different extraction solvents. *Food Res. Int.* 127,
575 108712.

576 Silva de Sá, I., Peron, A.P., Leimann, F.V., Bressan, G.N., Krum, B.N., Fachinetto, R., Pinela, J.,
577 Calhelha, R.C., Barreiro, M.F., Ferreira, I.C.F.R., Gonçalves, O.H., Ineu, R.P., 2019. In vitro
578 and in vivo evaluation of enzymatic and antioxidant activity, cytotoxicity and genotoxicity of
579 curcumin-loaded solid dispersions. *Food Chem. Toxicol.* 125, 29–37.

580 Soković, M., Glamočlija, J., Marin, P.D., Brkić, D., Griensven, L.J.L.D. van, 2010. Antibacterial
581 effects of the essential oils of commonly consumed medicinal herbs using an *in vitro* model.
582 *Molecules* 15, 7532–7546.

583 Spréa, R.M., Fernandes, Â., Calhelha, R.C., Pereira, C., Pires, T.C.S.P., Alves, M.J., Canan, C.,
584 Barros, L., Amaral, J.S., Ferreira, I.C.F.R., 2020. Chemical and bioactive characterization of
585 the aromatic plant: *Levisticum officinale* W.D.J. Koch: a comprehensive study. *Food Funct.*
586 11, 1292–1303.

587 Valdez-Solana, M.A., Mejía-García, V.Y., Téllez-Valencia, A., García-Arenas, G., Salas-Pacheco,
588 J., Alba-Romero, J.J., Sierra-Campos, E., 2015. Nutritional content and elemental and
589 phytochemical analyses of *Moringa oleifera* grown in Mexico. *J Chem.* 2015, 860381.

590 Ziani, B.E.C., Rached, W., Bachari, K., Alves, M.J., Calhelha, R.C., Barros, L., Ferreira, I.C.F.R.,
591 2019. Detailed chemical composition and functional properties of *Ammodaucus leucotrichus*
592 Cross. & Dur. and *Moringa oleifera* Lamarck. *J Funct Foods.* 53, 237–247.

593 Zungu, N., van Onselen, A., Kolanisi, U., Siwela, M., 2020. Assessing the nutritional composition
594 and consumer acceptability of *Moringa oleifera* leaf powder (MOLP)-based snacks for
595 improving food and nutrition security of children. *S. Afr. J. Bot.* 129, 283–290.

596

Figure captions

Fig. 1. *Moringa oleifera* leaf samples wild-harvested in Bolama (a and b) and commercially acquired, produced in Contuboel (c).

Table 1Nutritional value, free sugars and organic acids composition of *M. oleifera* leaves.

| | Wild sample | Commercial sample | <i>p</i> -value |
|--------------------------------|-------------|-------------------|-----------------|
| Nutritional value (g/100 g dw) | | | |
| Crude fat | 8.7±0.1 | 8.0±0.4 | 0.016 |
| Proteins | 27.4±0.3 | 32.6±0.9 | <0.001 |
| Ash | 12.4±0.1 | 8.63±0.09 | <0.001 |
| Total carbohydrates | 51.5±0.4 | 50.8±0.5 | 0.090 |
| Energy (kcal/100 g dw) | 393.7±0.2 | 406±2 | <0.001 |
| Free sugars (g/100 g dw) | | | |
| Fructose | 0.94±0.01 | 0.81±0.03 | 0.001 |
| Glucose | 0.41±0.01 | 0.57±0.03 | <0.001 |
| Sucrose | 1.97±0.02 | 5.33±0.09 | <0.001 |
| Trehalose | 0.84±0.01 | 1.20±0.05 | <0.001 |
| Total sugars | 4.16±0.02 | 7.9±0.2 | <0.001 |
| Organic acids (g/100 g dw) | | | |
| Oxalic acid | 1.83±0.01 | 1.97±0.01 | <0.001 |
| Malic acid | 3.95±0.04 | 4.08±0.01 | 0.001 |
| Ascorbic acid | 0.007±0.001 | 0.008±0.001 | 0.013 |
| Citric acid | 2.60±0.01 | 2.91±0.04 | <0.001 |
| Fumaric acid | tr | tr | - |
| Total organic acids | 8.38±0.04 | 8.97±0.06 | <0.001 |

Results are expressed mean ± SD ($n = 3$); tr: traces; Free sugars calibration curves: fructose ($y = 1.04x$, $R^2 = 0.999$; LOD = 0.05 mg/mL, LOQ = 0.18 mg/mL), glucose ($y = 0.935x$, $R^2 = 0.999$; LOD = 0.08 mg/mL; LOQ = 0.25 mg/mL); sucrose ($y = 0.977x$, $R^2 = 0.999$; LOD = 0.06 mg/mL, LOQ = 0.21 mg/mL) and trehalose ($y = 0.991x$, $R^2 = 0.999$; LOD = 0.07 mg/mL, LOQ = 0.24 mg/mL). Organic acids calibration curves: oxalic acid ($y = 9E + 10^6x + 459.731$; $R^2 = 0.994$; LOD = 12.55 µg/mL; LOQ = 41.82 µg/mL); malic acid ($y = 912.441x + 92.665$; $R^2 = 0.999$; LOD = 35.76 µg/mL; LOQ = 119.18 µg/mL); ascorbic acid ($y = 7E + 07x + 60.489$; $R^2 = 0.999$; LOD = 367 µg/mL; LOQ = 1222 µg/mL); citric acid ($y = 1E + 10^6x + 45.682$; $R^2 = 1$; LOD = 10.47 µg/mL; LOQ = 34.91 µg/mL) and fumaric acid ($y = 2E + 08x + 1E + 06$; $R^2 = 1$; LOD = 0.08 µg/mL; LOQ = 0.26 µg/mL).

Table 2Fatty acids and tocopherols composition of *M. oleifera* leaves.

| | Wild sample | Commercial sample | <i>p</i> -value |
|---------------------------|-------------|-------------------|-----------------|
| Fatty acids (relative %) | | | |
| C6:0 | 0.31±0.01 | 0.31±0.02 | 0.584 |
| C8:0 | 0.08±0.01 | 0.096±0.005 | 0.003 |
| C10:0 | 0.143±0.003 | 0.199±0.006 | <0.001 |
| C11:0 | 0.26±0.01 | 0.299±0.009 | 0.004 |
| C12:0 | 0.144±0.008 | 0.326±0.003 | <0.001 |
| C14:0 | 1.34±0.03 | 1.8±0.1 | 0.001 |
| C15:0 | 0.190±0.001 | 0.149±0.007 | <0.001 |
| C16:0 | 17.33±0.01 | 17.6±0.4 | 0.674 |
| C16:1 | 1.75±0.06 | 1.57±0.02 | 0.003 |
| C17:0 | 0.30±0.01 | 0.29±0.01 | 0.381 |
| C18:0 | 4.2±0.1 | 3.3±0.2 | <0.001 |
| C18:1n9c | 4.04±0.05 | 5.7±0.4 | <0.001 |
| C18:2n6c | 8.2±0.1 | 10.1±0.2 | <0.001 |
| C18:3n3 | 50.1±0.2 | 48.1±0.1 | <0.001 |
| C20:0 | 1.11±0.01 | 1.31±0.03 | <0.001 |
| C20:1 | 0.42±0.01 | 0.132±0.002 | <0.001 |
| C21:0 | 0.168±0.006 | 0.17±0.01 | 0.515 |
| C22:0 | 4.1±0.1 | 4.0±0.3 | 0.359 |
| C22:2 | 0.48±0.03 | 0.246±0.005 | <0.001 |
| C23:0 | 0.40±0.01 | 0.28±0.01 | <0.001 |
| C24:0 | 4.73±0.09 | 4.07±0.09 | <0.001 |
| SFA | 35.0±0.3 | 34.1±0.5 | 0.019 |
| MUFA | 6.21±0.02 | 7.4±0.4 | 0.001 |
| PUFA | 58.8±0.3 | 58.5±0.1 | 0.75 |
| Tocopherols (mg/100 g dw) | | | |
| α-Tocopherol | 58.79±0.03 | 68.03±0.05 | <0.001 |
| β-Tocopherol | 3.23±0.04 | 2.41±0.06 | <0.001 |
| δ-Tocopherol | 10.85±0.07 | 6.63±0.01 | <0.001 |
| Total tocopherols | 72.87±0.06 | 77.1±0.1 | <0.001 |

Results are expressed mean ± SD (*n* = 3); caproic acid (C6:0), caprylic acid (C8:0), capric acid (C10:0), undecanoic acid (C11:0), lauric acid (C12:0), myristic acid (C14:0), pentadecanoic acid (C15:0), palmitic acid (C16:0), palmitoleic acid (C16:1), heptadecanoic acid (C17:0), stearic acid (C18:0), oleic acid (C18:1n9c), linoleic acid (C18:2n6c), α-linolenic acid (C18:3n3), arachidic acid (C20:0), *cis*-11-eicosenoic acid (C20:1), eicosenoic acid (C21:0), behenic acid (C22:0), *cis*-13,16-docosadienoic acid (C22:2), tricosanoic acid (C23:0), and lignoceric acid (C24:0); SFA: saturated fatty acids; MUFA: monounsaturated fatty acids; PUFA: polyunsaturated fatty acids; tr: traces. Tocopherols

calibration curves: α -tocopherol ($y = 1.295x$; $R^2 = 0.991$; LOD: 18.06 ng/mL, LOQ: 60.20 ng/mL); β -tocopherol ($y=0.396x$; $R^2 = 0.992$; LOD: 25.82 ng/mL, LOQ: 86.07 ng/mL); γ -tocopherol ($y = 0.567x$; $R^2 = 0.991$; LOD: 14.79 ng/mL, LOQ: 49.32 ng/mL); δ -tocopherol ($y = 0.678x$; $R^2 = 0.992$; LOD: 20.09 ng/mL, LOQ: 66.95 ng/mL).

Table 3

Phenolic compounds identified in hydroethanolic, infused and decocted *M. oleifera* leaf extracts. It is presented the retention time (Rt), wavelengths of maximum absorption in the visible region (λ_{\max}), and mass spectral data.

| Peak | Rt (min) | λ_{\max} (nm) | [M-H] (m/z) | MS ² (m/z) | Tentative identification |
|------|----------|-----------------------|-------------|--|---|
| 1 | 4.15 | 324 | 341 | 179(100), 161(3), 135(40) | Caffeic acid hexoside |
| 2 | 4.69 | 324 | 353 | 191(100), 179(47), 173(5), 161(3), 135(10) | 3- <i>O</i> -Caffeoylquinic acid |
| 3 | 4.88 | 324 | 353 | 191(20), 179(55), 173(100), 161(5), 135(8) | 4- <i>O</i> -Caffeoylquinic acid |
| 4 | 6.25 | 280 | 337 | 191(8), 173(5), 163(100), 155(5), 137(3), 119(5) | 3- <i>p</i> -Coumaroylquinic acid |
| 5 | 6.40 | 324 | 353 | 191(100), 179(80), 173(24), 161(5), 135(10) | <i>cis</i> -5- <i>O</i> -Caffeoylquinic acid |
| 6 | 7.08 | 324 | 353 | 191(100), 179(80), 173(42), 161(5), 135(12) | <i>trans</i> -5- <i>O</i> -Caffeoylquinic acid |
| 7 | 9.86 | 338 | 593 | 473(100), 383(15), 353(25) | Apigenin-6,8-di- <i>C</i> -hexoside |
| 8 | 10.29 | 338 | 593 | 473(100), 383(15), 353(25) | Apigenin 8- <i>C</i> -glucoside 6"- <i>O</i> -glucoside |
| 9 | 11.39 | 338 | 593 | 473(100), 383(15), 353(25) | Apigenin 6- <i>C</i> -glucoside 7- <i>O</i> -glucoside |
| 10 | 13.47 | 338 | 593 | 473(100), 383(15), 353(25) | Apigenin 6,8- <i>C</i> -di- β - <i>D</i> -glucopyranoside |
| 11 | 14.86 | 327 | 447 | 301(100) | Quercetin-3- <i>O</i> -rhamnoside |
| 12 | 16.43 | 352 | 431 | 341(27), 311(100) | Apigenin-8- <i>C</i> -glucoside |
| 13 | 17.67 | 352 | 609 | 301(100) | Quercetin-3- <i>O</i> -rutinoside |
| 14 | 18.24 | 352 | 609 | 301(100) | Quercetin-4'- <i>O</i> -rhamnoside-7- <i>O</i> -hexoside |
| 15 | 18.86 | 352 | 609 | 301(100) | Quercetin-3- <i>O</i> -rhamnoside-7- <i>O</i> -hexoside |
| 16 | 19.02 | 352 | 463 | 301(100) | Quercetin 3- <i>O</i> -glucoside |
| 17 | 19.77 | 350 | 505 | 301(100) | Quercetin-3- <i>O</i> -(6''-acetyl)-glucoside |
| 18 | 20.11 | 350 | 549 | 505(10), 463(27), 301(100) | Quercetin-3- <i>O</i> -(6''-malonyl)-glucoside) |
| 19 | 21.00 | 350 | 549 | 505(10), 463(27), 301(100) | Quercetin 7- <i>O</i> -malonylglucoside |
| 20 | 22.04 | 350 | 549 | 505(10), 463(27), 301(100) | Quercetin- <i>O</i> -malonyl-hexoside |
| 21 | 23.4 | 350 | 549 | 505(10), 463(27), 301(100) | Quercetin 3- <i>O</i> -malonylglucoside |
| 22 | 24.07 | 350 | 623 | 315(100) | Isorhamnetin-3- <i>O</i> -rutinoside |
| 23 | 24.59 | 343 | 533 | 285(100) | Kaempferol-3- <i>O</i> -(6''-malonyl)-glucoside) |
| 24 | 25.33 | 349 | 519 | 315(100) | Isorhamnetin-3- <i>O</i> -(6''-acetyl)-glucoside |
| 25 | 25.88 | 350 | 563 | 315(100) | Isorhamnetin-3- <i>O</i> -(6''-malonyl)-glucoside) |

Table 4Content (mg/g of extract) of the phenolic compounds identified in the hydroethanolic, infused and decocted *M. oleifera* leaf extracts.

| Peak | Wild sample extracts | | | Commercial sample extracts | | |
|------------|-------------------------|-------------------------|------------------------|----------------------------|------------------------|------------------------|
| | Hydroethanolic | Infused | Decocted | Hydroethanolic | Infused | Decocted |
| 1 | 2.6±0.1 | nd | nd | nd | nd | nd |
| 2 | 5.5±0.1 ^e | 4.59±0.01 ^d | 6.07±0.09 ^b | 7.4±0.1 ^a | 5.8±0.4 ^c | 5.9±0.3 ^c |
| 3 | 5.49±0.05 ^d | 2.06±0.02 ^e | nd | 5.9±0.2 ^c | 6.2±0.3 ^b | 12.1±0.3 ^a |
| 4 | 1.31±0.03 ^c | nd | 2.21±0.08 ^b | 1.23±0.03 ^d | 2.97±0.04 ^a | nd |
| 5 | 5.6±0.2 ^c | 5.75±0.09 ^c | 4.7±0.3 ^d | 4.1±0.3 ^e | 6.16±0.08 ^b | 8.3±0.4 ^a |
| 6 | nd | nd | nd | 2.51±0.08 ^b | nd | 7.3±0.4 ^a |
| 7 | 8.38±0.41 ^b | 9.9±0.5 ^a | nd | 3.3±0.1 ^e | 5.8±0.1 ^c | 4.25±0.04 ^d |
| 8 | 13.8±0.1 ^a | 3.98±0.04 ^c | 9.5±0.4 ^b | 2.7±0.1 ^d | nd | 8.7±0.2 ^c |
| 9 | 7.49±0.05 ^a | nd | 4.1±0.2 ^c | 3.47±0.05 ^d | nd | 4.6±0.2 ^b |
| 10 | 4.98±0.20 | nd | nd | nd | nd | nd |
| 11 | 1.26±0.07 | nd | nd | nd | nd | nd |
| 12 | 6.27±0.05 ^a | 2.4±0.1 ^b | nd | 1.7±0.1 ^c | nd | nd |
| 13 | 13.7±0.3 ^a | 6.94±0.08 ^e | 10.5±0.5 ^c | 11.6±0.5 ^b | 8.0±0.3 ^d | 7.78±0.07 ^d |
| 14 | 5.4±0.3 ^a | 4.2±0.2 ^b | 3.33±0.07 ^d | 3.9±0.2 ^c | 5.33±0.08 ^a | 3.95±0.02 ^c |
| 15 | 7.1±0.2 ^a | 3.1±0.1 ^b | 1.27±0.07 ^d | nd | 1.91±0.03 ^c | nd |
| 16 | nd | nd | nd | 3.2±0.1 | nd | nd |
| 17 | nd | 3.6±0.2 ^d | nd | 7.9±0.9 ^a | 6.3±0.2 ^b | 4.20±0.09 ^c |
| 18 | 6.7±0.2 ^a | 1.59±0.06 ^d | 2.3±0.1 ^c | 2.62±0.05 ^b | 1.66±0.09 ^d | 2.42±0.03 ^c |
| 19 | 2.8±0.1 ^a | 1.36±0.05 ^c | 1.61±0.04 ^b | 1.26±0.06 ^d | nd | nd |
| 20 | 1.66±0.09 ^a | nd | nd | 0.46±0.01 ^b | nd | nd |
| 21 | 1.23±0.02 ^a | nd | nd | 1.09±0.05 ^b | nd | 0.46±0.01 ^c |
| 22 | nd | nd | nd | nd | nd | 1.39±0.05 |
| 23 | 1.41±0.03 ^a | 1.11±0.03 ^c | 0.81±0.02 ^e | nd | 1.23±0.03 ^b | 1.03±0.02 ^d |
| 24 | nd | nd | nd | nd | 0.94±0.02 | nd |
| 25 | 1.11±0.05 ^a | 1.07±0.05 ^b | 0.91±0.02 ^c | 0.93±0.04 ^c | nd | nd |
| TPA | 20.6±0.3 ^c | 12.41±0.08 ^e | 13±1 ^d | 21.1±0.1 ^b | 21.1±0.5 ^b | 33.5±0.7 ^a |
| TF | 83.15±0.07 ^a | 39.4±0.6 ^c | 34±1 ^e | 44.2±0.4 ^b | 31.1±0.5 ^f | 38.7±0.3 ^d |
| TPC | 103.7±0.2 ^a | 51.8±0.5 ^d | 47±1 ^e | 65.3±0.3 ^c | 52±1 ^d | 72±1 ^b |

Results are expressed mean ± SD ($n = 3$); nd: not detected; TPA: total phenolic acids; TF: total flavonoids; TPC: total phenolic compounds. Different letters in the same line indicate significant differences according to a Tukey's HSD test ($p < 0.05$). The peak identification is provided in **Table 3**. Calibration curves used in the quantification: Standard calibration curves: caffeic acid ($y = 388345x + 406369$, $R^2 = 0.999$, limit of detection (LOD) = 0.78 µg/mL and limit of quantitation

(LOQ) = 1.97 $\mu\text{g/mL}$, peaks 1); chlorogenic acid ($y = 168823x - 161172$, $R^2 = 0.999$, LOD = 0.20 $\mu\text{g/mL}$ and LOQ = 0.68 $\mu\text{g/mL}$, peaks 2, 3, 5 and 6); *p*-coumaric acid ($y = 301950x + 6966.7$, $R^2 = 0.9999$, LOD = 0.68 $\mu\text{g/mL}$ and LOQ = 1.61 $\mu\text{g/mL}$, peaks 4); apigenin-6-*C*-glucoside ($y = 107025x + 61531$, $R^2 = 0.9989$, LOD = 0.19 $\mu\text{g/mL}$ and LOQ = 0.63 $\mu\text{g/mL}$, peaks 7, 8, 9, 10 and 12); quercetin-3-*O*-glucoside ($y = 34843x - 160173$, $R^2 = 0.9998$, LOD = 0.21 $\mu\text{g/mL}$ and LOQ = 0.71 $\mu\text{g/mL}$, peaks 11, 16, 17, 18, 19, 20, 21, 22, 23, 24 and 25); quercetin-3-*O*-glucoside ($y = 13343x + 76751$, $R^2 = 0.9998$, LOD = 0.21 $\mu\text{g/mL}$ and LOQ = 0.71 $\mu\text{g/mL}$, peaks 13, 14 and 15).

Table 5Antioxidant activity of the hydroethanolic, infused and decocted *M. oleifera* leaf extracts.

| Antioxidant assay* | Extract | Wild sample | Commercial sample | <i>p</i> -value |
|----------------------------|----------------|-------------|-------------------|-----------------|
| TBARS | Hydroethanolic | 54±3 | 57±4 | 0.449 |
| | Infused | 318±6 | 232±4 | <0.001 |
| | Decocted | 486±8 | 417±6 | <0.001 |
| OxHLIA, Δ <i>t</i> 60 min | Hydroethanolic | 70±2 | 81±5 | <0.001 |
| | Infused | 13±1 | 1.8±0.2 | <0.001 |
| | Decocted | 14.5±0.2 | 47±2 | <0.001 |
| OxHLIA, Δ <i>t</i> 120 min | Hydroethanolic | na | na | - |
| | Infused | 30±2 | 6.1±0.2 | <0.001 |
| | Decocted | 49±2 | 104±3 | <0.001 |

*Results are expressed as IC₅₀ values (μg/mL); na: no activity; Trolox IC₅₀ values: 23±0.1 μg/mL (TBARS), 19.6±0.8 μg/mL (OxHLIA, Δ*t* 60 min) and 41±1 μg/mL (Δ*t* 120 min).

Table 6

Antibacterial (minimal inhibitory and minimal bactericidal concentrations (MIC and MBC, respectively) in mg/mL) and antifungal (MIC and minimal fungicidal concentration (MFC) in mg/mL) activity of the hydroethanolic, infused and decocted *M. oleifera* leaf extracts.

| Microorganisms | Extract | Wild sample | Commercial sample | Streptomycin | Ampicillin |
|-------------------------|----------------|-------------|-------------------|--------------|------------|
| | | | | MIC/MBC | MIC/MBC |
| Bacterial strains | | MIC/MBC | MIC/MBC | MIC/MBC | MIC/MBC |
| <i>B. cereus</i> | Hydroethanolic | 0.037/0.15 | 0.075/0.15 | | |
| | Infused | 0.037/0.15 | 0.15/0.3 | 0.04/0.1 | 0.25/0.45 |
| | Decocted | 0.15/0.3 | 0.075/0.15 | | |
| <i>S. aureus</i> | Hydroethanolic | 0.1/0.15 | 0.15/0.3 | | |
| | Infused | 0.1/0.15 | 0.6/0.9 | 0.1/0.2 | 0.25/0.4 |
| | Decocted | 0.3/0.60 | 0.15/0.3 | | |
| <i>L. monocytogenes</i> | Hydroethanolic | 0.1/0.15 | 0.15/0.3 | | |
| | Infused | 0.1/0.15 | 0.15/0.3 | 0.2/0.3 | 0.4/0.5 |
| | Decocted | 0.15/0.6 | 0.15/0.3 | | |
| <i>E. coli</i> | Hydroethanolic | 0.05/0.075 | 0.075/0.15 | | |
| | Infused | 0.05/0.075 | 0.075/0.15 | 0.2/0.3 | 0.4/0.5 |
| | Decocted | 0.15/0.3 | 0.15/0.3 | | |
| <i>E. cloacae</i> | Hydroethanolic | 0.15/0.3 | 0.3/0.6 | | |
| | Infused | 0.1/0.3 | 0.6/0.9 | 0.2/0.3 | 0.25/0.5 |
| | Decocted | 0.15/0.3 | 0.3/0.6 | | |
| <i>S. typhimurium</i> | Hydroethanolic | 0.1/0.15 | 0.15/0.3 | | |
| | Infused | 0.1/0.15 | 0.2/0.3 | 0.2/0.3 | 0.75/1.2 |
| | Decocted | 0.15/0.3 | 0.15/0.3 | | |
| Fungal strains | | MIC/MFC | MIC/ MFC | Ketoconazole | Bifonazole |
| | | | | MIC/ MFC | MIC/ MFC |
| <i>A. fumigatus</i> | Hydroethanolic | 0.075/0.15 | 0.075/0.15 | | |
| | Infused | 0.3/0.6 | 0.90/1.2 | 0.25/0.5 | 0.15/0.2 |
| | Decocted | 0.075/0.15 | 0.075/0.15 | | |
| <i>A. ochraceus</i> | Hydroethanolic | 0.037/0.075 | 0.037/0.075 | | |
| | Infused | 0.037/0.075 | 0.15/0.3 | 0.2/0.5 | 0.10/0.2 |
| | Decocted | 0.037/0.075 | 0.037/0.075 | | |
| <i>A. niger</i> | Hydroethanolic | 0.075/0.15 | 0.037/0.075 | | |
| | Infused | 0.075/0.15 | 0.075/0.15 | 0.2/0.5 | 0.15/0.2 |
| | Decocted | 0.037/0.075 | 0.02/0.037 | | |
| <i>P. funiculosum</i> | Hydroethanolic | 0.05/0.075 | 0.1/0.2 | | |
| | Infused | 0.05/0.075 | 0.6/0.9 | 0.20/0.50 | 0.20/0.25 |
| | Decocted | 0.075/0.15 | 0.075/0.15 | | |
| <i>P. ochrochloron</i> | Hydroethanolic | 0.15/0.3 | 0.075/0.3 | | |
| | Infused | 0.1/0.15 | 0.6/1.2 | 2.5/3.5 | 0.20/0.25 |
| | Decocted | 0.15/0.3 | 0.075/0.3 | | |

| | | | | | |
|-----------------------|----------------|------------|-----------|---------|---------|
| <i>P.v. cyclopium</i> | Hydroethanolic | 0.075/0.15 | 0.075/0.3 | | |
| | Infused | 0.075/0.15 | 0.6/0.9 | 0.2/0.3 | 0.1/0.2 |
| | Decocted | 0.075/0.15 | 0.075/0.3 | | |
