

Associations between syndesmophytes and facet joint ankylosis in radiographic axial spondyloarthritis patients on low-dose CT over 2 years

Rosalinde Stal¹, Alexandre Sepriano^{1,2}, Floris Alexander van Gaalen¹, Xenofon Baraliakos³, Rosaline van den Berg¹, Monique Reijnen¹, Juergen Braun³, Robert B M Landewé^{4,5}, Désirée van der Heijde¹

¹Leiden University Medical Center, Leiden, the Netherlands, ²Universidade Nova de Lisboa, NOVA Medical School, Lisbon, Portugal, ³Ruhr-University Bochum, Rheumazentrum Ruhrgebiet, Herne, Germany, ⁴Amsterdam UMC locatie AMC, Amsterdam, the Netherlands, ⁵Zuyderland Medical Centre Heerlen, Heerlen, the Netherlands

Corresponding author

Rosalinde Stal, ORCID iD (0000-0002-6706-3933)

Stafcentrum Reumatologie

Albinusdreef 2

2333ZA Leiden

The Netherlands

Keywords

radiographic axial spondyloarthritis; syndesmophytes; facet joints; ankylosis; low dose CT

Rheumatology key messages

- Bridging syndesmophytes increase risk of facet joint ankylosis in the same and adjacent vertebral units
- Facet joint ankylosis increases risk of syndesmophytes in the same and adjacent vertebral units
- Inflammation in posterior elements was positively associated with facet joint ankylosis two years later

Abstract

Objectives. In radiographic axial spondyloarthritis (r-axSpA), spinal damage manifests as syndesmophytes and facet joint ankylosis (FJA). We evaluated whether the presence of one lesion increased the risk of the other lesion.

Methods. Patients with r-axSpA underwent low-dose CT (IdCT) and MRI of the whole spine at baseline and two years. On IdCT, vertebrae were scored for presence and size of syndesmophytes; facet joints were assessed for ankylosis. MR images were assessed for inflammation. Two hypotheses were tested: 1) presence of FJA is associated with new syndesmophyte(s) on the same vertebral unit (VU) two years later, and 2) presence of bridging syndesmophyte(s) is associated with new FJA on the same VU two years later. Two Generalized Estimating Equations models were tested per hypothesis using increase of FJA/syndesmophytes (model A) or presence of FJA/syndesmophytes (model B) as outcome, adjusted for inflammation at baseline. Secondary analyses tested the hypotheses with outcomes on adjacent VUs and dose-response effects.

Results. Fifty-one patients were included (mean age 49, 84% male, 82% HLA-B27+). Baseline bridging syndesmophytes occurred more often (range: 10-60% per VU) than FJA (range: 8-36%). ORs(95%CI) for presence of bridging syndesmophytes on development of FJA were 3.55(2.03-6.21) for model A and 3.30(2.14-5.09) for model B. ORs for presence of baseline FJA on new syndesmophytes were 1.87(1.20-2.92) for model A and 1.69(0.88-3.22) for model B. Secondary analyses yielded positive ORs for both hypotheses.

Conclusions. Bone formation in vertebrae and in facet joints influence each other's occurrence, with the effect of syndesmophytes being larger than that of FJA.

Introduction

Radiographic axial spondyloarthritis (r-axSpA) is characterized by inflammation and structural damage in the sacroiliac joints and, in a subset of patients, the spine.[1] Spinal lesions can occur in several locations, including the vertebrae, facet joints and ligaments.[2] Syndesmophytes are osseous spikes on the vertebral rim, being formed from ossification of the annulus fibrosis, anterior longitudinal ligament and paravertebral connective tissue, and growing in the direction of the adjacent vertebra.[3] Fusion of two vertebrae occurs when a syndesmophyte has crossed the intervertebral disc space and formed a so-called bridging syndesmophyte to the adjacent vertebra.[4] Pathological bone formation has also been shown to occur in the facet joints (also known as (zyg)apophyseal joints), causing facet joint ankylosis (FJA).[5, 6] There is conflicting data on the direction of the association between FJA and (bridging) syndesmophytes. One study reported that bridging syndesmophytes occur infrequently without FJA, hence proposing that facet joints are primarily involved in r-axSpA.[7] Another study reported that FJA was stronger associated with bridging than non-bridging syndesmophytes and that thoracolumbar syndesmophytes occurred often without FJA, therefore suggesting that syndesmophyte development precedes FJA.[8] Several techniques have been used over the years to study structural lesions in the spine, with lateral cervical and lumbar conventional radiography (CR) being the most widely used technique.[9-12] In recent years, low dose Computed Tomography (IdCT) has been shown to be an adequate and feasible technique for repeated imaging of the whole spine.[13, 14] Its ability to capture the thoracic spine gives it certain superiority over CR as an imaging tool, especially since syndesmophytes have been reported to occur and progress most in the thoracic spine.[9, 15] Furthermore, its drastically lowered radiation dose relative to conventional CT makes it a feasible technique to apply repeatedly to the whole spine.

Multiple studies have looked into factors associated with syndesmophyte development. Associations have been made with factors including disease activity, disease duration, smoking, age, gender and presence of existing syndesmophytes, the latter being the most prominent and frequently confirmed contributing factor.[16] Other studies, looking more into the pathophysiological process, found significant effects of vertebral corner inflammation and vertebral corner fat deposition on the development of syndesmophytes.[17, 18] Since we know that presence of existing syndesmophytes

is a predictor for further syndesmophyte development, the question arises if this is due to the rigidity that is caused by the syndesmophytes, and perhaps rigidity elsewhere in the spinal column. In the current study we hypothesize a mechanical effect of rigidity in the spine resulting in bone formation, looking specifically at the interplay between FJA and syndesmophytes in the same area of the spine. Using whole spine IdCT we study whether presence of bridging syndesmophytes increases the risk of FJA, and whether presence of FJA increases the risk of syndesmophyte formation at the same or adjacent vertebral level.

Methods

Patients

Data were used from the Sensitive Imaging in Ankylosing Spondylitis (SIAS) cohort, which included r-axSpA patients from Leiden, the Netherlands, and Herne, Germany. The study was approved by the medical ethical committees of Leiden (Medisch Ethische Toetsings Commissie, P10.021) and Herne (Ethikkommission der Ruhr Universität Bochum, 4366-12). Patients fulfilled the modified New York criteria, had at least one inflammatory lesion on spinal Magnetic Resonance Imaging (MRI) and between one and 18 syndesmophytes on lateral cervical and lumbar conventional radiography (CR). All patients gave written informed consent.

Imaging techniques

LdCT and MR images of the whole spine were collected at baseline and two years. MR images with 3.5mm sagittal slices were obtained on a 3Tesla (Leiden) and 1.5Tesla scanner (Leiden: Philips Medical Systems, Best, The Netherlands. Herne: Siemens Aera 1.5, Siemens, Erlangen, Germany). LdCT images with 1mm axial slices and 2mm sagittal and coronal slices were obtained on a 64-section (Leiden) and 16-section CT scanner (Leiden: Aquilion 64, Toshiba Medical Systems, Otawara, Japan. Herne: Somatom Emotion 16, Siemens, Erlangen, Germany).

IdCT scoring methods and variable definitions

IdCT mages were assessed for presence and size of syndesmophytes on sagittal and coronal slices with the Computed Tomography Syndesmophyte Score (CTSS) and for presence of FJA on axial slices by two trained, central readers. The CTSS has been described in detail in a separate publication.[14] In short, the CTSS assesses four quadrants of a vertebral unit (VU) per plane. A VU comprises the lower half of a vertebra, the upper half of the vertebra underneath and the intervertebral disc space (IDS) in between (Supplementary Figure S1, available at *Rheumatology* online). Scores use a four-point scale per quadrant: 0) no syndesmophyte; 1) syndesmophyte reaching <50% of the IDS; 2) syndesmophyte reaching \geq 50% of the IDS; 3) syndesmophyte bridging the IDS. Thus, a maximum of eight syndesmophytes or four bridging syndesmophytes can be scored with the CTSS per VU. For both syndesmophyte and FJA scoring, the readers could attribute a missing score if the location was difficult to assess due to e.g. image quality.

Dichotomous status scores at baseline and follow-up were made per reader, per VU, to show whether there was presence of at least one (out of eight) syndesmophyte. Furthermore, a change score was made per VU, per reader, reflecting whether there was an increase in the number of syndesmophytes over time. The change score was set to missing if all non-missing quadrants in a VU had a syndesmophyte at baseline and thus could not show change over time. A status score at baseline was made to show whether there was presence of at least one bridging syndesmophyte in the VU. Lastly, a status score at baseline was made showing how many quadrants of the VU had a bridging syndesmophyte (range 0-4). Presence of FJA was coded dichotomously per reader for the left and right facet joint. For FJA, similar variables were made as described above. Dichotomous, individual reader status scores at baseline and follow-up showed whether there was presence of at least one (out of two) ankylosed facet joint per VU. A change score showed whether there was an increase in the number of ankylosed facet joints over time, and this was set to missing if all facet

1
2
3 joints with non-missing scores in a VU were ankylosed at baseline. Lastly, a status score at baseline
4 was made showing how many facet joints were ankylosed (range 0-2).
5

6 **MRI scoring methods and variable definitions**

7 Bone marrow oedema suggestive of spondyloarthritis was scored by three trained central readers on
8 short-tau inversion recovery (STIR) images. Inflammation on the vertebral bodies was scored with
9 the Spondyloarthritis Research Consortium of Canada (SPARCC) scoring system and coded as
10 inflammation present or absent per VU based on agreement by $\geq 2/3$ readers. [19] Inflammation in
11 the posterior elements (pedicles and soft tissue in C2-T1; facet joints, processes, pedicles and soft
12 tissues in T1-S1) was also coded as present or absent per VU if inflammation was present in at least
13 one of the posterior elements according to $\geq 2/3$ readers.
14
15

16 **Statistical analyses**

17 To test whether there is a longitudinal relationship between syndesmophytes and FJA, two
18 hypotheses were formulated, looking at both directions: 1) presence of FJA is associated with a new
19 syndesmophyte on the same VU two years later, and 2) presence of a bridging syndesmophyte is
20 associated with new FJA on the same VU two years later. We used multivariable multilevel
21 Generalized Estimating Equation (GEE) models to assess the hypotheses at the VU-level and to use
22 scores from each individual reader, which increases statistical power. [20] Because the models take
23 into account correlations within patients (the correlation between VUs from the same patient) next
24 to the change in VUs on different timepoints, the effects should be interpreted as truly longitudinal.
25 An exchangeable working correlation structure was used to handle the VU-level.
26
27

28 Each hypothesis was tested in two types of models. Model A ('change-score' model) looked at the
29 effect of the predictor at baseline on the increase in the number of lesions at follow-up. For
30 example, for hypothesis 1, model A studied the effect of the presence of FJA at baseline on the
31 syndesmophyte change score (≥ 1 new syndesmophyte). Model B ('autoregressive' model) looked at
32 the effect of the predictor at baseline on the presence of a lesion in a VU at follow-up adjusting for
33 its presence at baseline (the 'autoregressor'). For example, for hypothesis 2, model B studied the
34 effect of a bridged syndesmophyte at baseline on presence of ≥ 1 ankylosed facet joint in a VU at
35 follow-up (status score), adjusted for the presence of ≥ 1 ankylosed facet joint in the VU at baseline.
36 All models were adjusted for the presence of inflammation at baseline on the location of the
37 outcome, e.g., presence of inflammation at baseline on the vertebral body for hypothesis 1 and
38 presence of inflammation in the posterior elements at baseline for hypothesis 2.
39
40

41 Additional analyses were performed (using also model A and B), to 1) assess the hypotheses on
42 adjacent VUs and, 2) assess a dose-response effect between the bony lesions. For additional
43 analyses 1 associations were tested between the predictor on one VU and the outcome on the VU
44 above, the VU below, two VUs above and two VUs below (see Supplementary Figure S2, available at
45 *Rheumatology* online). The models were again adjusted for inflammation at baseline on the location
46 of the outcome. For additional analyses 2 the same models as for the primary analyses were used,
47 but now replacing the binary predictor (for FJA present/absent at baseline and bridging
48 syndesmophyte(s) present/absent at baseline) with the categorical predictor (ankylosis in 0, 1 or 2
49 facet joints and bridging syndesmophytes in 0, 1, 2, 3 or 4 quadrants). These models are used to
50 assess presence of a dose-response effect, ie. does the risk of developing the outcome increase if
51 there is more presence of the predictor. All previously mentioned analyses were performed on a
52 group level. To assess the relationship between the presence of bridging syndesmophytes and FJA
53 on a patient level a cumulative probability plot was made of the average reader scores per patient of
54 the number of VUs with at least one bridging syndesmophyte and the number of VUs with at least
55 one ankylosed facet joint. This figure is used to assess the frequency of occurrence of both lesions
56 within each patient, to see which occurs more often.
57
58
59
60

Intraclass Correlations Coefficients (ICCs) for syndesmophyte scores and FJA scores were previously published.[6, 14] The ICC for inflammation in the posterior elements was calculated on the patient level for baseline status scores using two-way mixed effect average ICCs, and was 0.82. The ICC for inflammation on the vertebral bodies at baseline, with the same model, was 0.89.

Previously, the ability of IdCT to assess FJA was evaluated, and VUs from C5-C6 until T1-T2 were deemed difficult to assess.[6] Therefore, facet joint scores from these four VUs were excluded from all analyses.

Patient and public involvement

There is structural patient participation in all research projects at the Leiden University Medical Center's department of Rheumatology. This is achieved through a patient council. The current study is in line with the council's wish to prevent development and progression of structural damage in r-axSpA. Patients were not directly involved in designing or conducting the study, but patient(s) were part of the medical ethical committee for both centers.

Results

LdCT scans at baseline and two years and MRI scans at baseline from 51 patients were included in the analyses (mean age 49 years (SD 10), 68% male, 78% HLA-B27+)(Supplementary Table S1, available at *Rheumatology* online). The occurrence of inflammation is presented in Supplementary Figure S2. The presence of bridging syndesmophytes and FJA at baseline per reader per VU is presented in figure 1. Both FJA and bridging syndesmophytes are present at baseline in every VU in at least one patient. Both lesions occur more often in the thoracic spine, bridging syndesmophytes occur more often than FJA at essentially all levels. Figure 2 presents this data on the patient level, showing that in the vast majority of all patients, the number of VUs with bridging syndesmophytes exceeds the number of VUs with FJA.

Per patient, 23 VUs were imaged per timepoint, yielding a total of 1173 VUs to be assessed over time. After excluding FJA scores for C5-C6 until T1-T2, missing scores for IdCT images due to e.g. poor visibility were low, ranging from 0.5% to 2% for CTSS and 1.9% to 4.4% for FJA scores on baseline and follow-up for both readers. VUs with missing scores were not imputed and were excluded from analyses.

For reader 1, 999 VUs had no syndesmophytes at baseline (N=451), or had a syndesmophyte on ≥ 1 but not all quadrants of the VU (N=548). Of these, 16% (164/999) developed a syndesmophyte after 2 years. Of the 451 VUs that had no syndesmophyte at baseline, 10% (44/451) developed a syndesmophyte. Reader 2 reported similar results with 18% (186/1008) and 11% (45/410). For reader 1, 750 VUs had zero (N=728) or one (N=22) ankylosed facet joint at baseline. Of these, 5% (41/750) of VUs developed FJA after 2 years. Of the 728 VUs without FJA at baseline, 5% (36/728) developed FJA. Reader 2 reported similar results of 7% (52/718) and 6% (42/680).

Main model results

Results for the main analyses are presented in table 1. Odds ratios (ORs) for the association between FJA at baseline and syndesmophyte development at follow-up (hypothesis 1) were 1.87 (1.20-2.92) for the increase in the number of syndesmophytes (model A), and 1.69 (0.88-3.22) for VUs with new presence of syndesmophytes (model B). In both models, vertebral body inflammation at baseline was significantly associated with the outcome. ORs for the association between bridging syndesmophytes at baseline and FJA development at follow-up (hypothesis 2) were slightly higher and significant for both models, with 3.55 (2.03-6.21) for model A and 3.30 (2.14-5.09) for model B. In these models, the association between inflammation in the posterior elements at baseline was only significant for model B.

Additional model results

Results for the analyses on adjacent VUs are presented in table 2. For hypothesis 1, statistically significant ORs for the association between FJA at baseline and syndesmophyte development at follow-up were found on one VU above or below and two VUs below. For hypothesis 2, significant ORs for the association between bridging syndesmophytes at baseline and FJA development at follow-up were also found for one VU above or below and two VUs below.

Results for the analyses assessing presence of a possible dose-response effect are presented in table 3. For all models there is an upwards trend where the odds ratios become larger when there is more presence of the predictor. For models including FJA as a predictor effects remain small, the greatest being the effect of having two ankylosed facet joints on the increase of syndesmophytes (2.15(1.24-3.27)). For models including bridging syndesmophytes as a predictor the effects increase drastically: having one quadrant (compared to zero quadrants) with a bridged syndesmophyte yields an OR of 2.22 (0.90-5.46) for the increase of FJA and 2.35 (1.27-4.34) for the presence of FJA, while having four quadrants with bridged syndesmophytes (compared to zero quadrants) yields an OR of 6.43 (2.95-14.03) for the increase of FJA and 5.54 (3.14-9.80) for the presence of FJA.

Discussion

In the current study we assessed whether FJA was associated with syndesmophyte development two years later (hypothesis 1), and whether fusion of the vertebral bodies in the form of bridging syndesmophytes was associated with FJA (hypothesis 2), with both associations being adjusted for presence of inflammation. We present evidence supporting both hypotheses, with the strongest associations for bridging syndesmophytes being a predictor for FJA. We also found evidence for presence of a dose-response effect in both directions with high odds ratios when there was extensive presence of bridging syndesmophyte, further supporting the hypotheses. In addition to studying the hypotheses on the same VU, we explored the hypotheses with the outcome on one or two VUs above or below the predictor. We found statistically significant positive associations for five out of eight models for both hypotheses, indicating that FJA or bridging syndesmophytes in an adjacent VU increase risk of syndesmophyte development or FJA. Just as for the main models, ORs are slightly higher for bridging syndesmophytes being a predictor for FJA (hypothesis 2) than for the FJA being a predictor for syndesmophyte development (hypothesis 1).

The literature provided little, and somewhat contradicting, information on associations between FJA and syndesmophytes. As we found both syndesmophytes and FJA to be risk factors for subsequent bone formation, our results do not support the notion that FJA is a primary lesion in r-axSpA.[7] On the contrary, syndesmophytes appeared to be stronger risk factors for FJA than vice versa. This is also substantiated by the fact that bridging syndesmophytes are more common than FJA in our study population. With these findings, our study is more in line with the study by Tan et al.,[8] reporting that syndesmophytes are likely to appear before FJA on a vertebral level. It is important to note, however, that our study population was selected for the presence of at least one syndesmophyte and one inflammatory spinal lesion at baseline, and therefore the results need to be interpreted in this context. Our sample size was also modest (N=51), although the use of IdCT allowed to assess the whole spine of each patient thus yielding a 'sample size' of 1173 VUs. Our imaging assessments were performed twice and with a two-year interval. It is possible that a longer time interval is needed to fully capture the effects under study. Therefore, future studies are needed to confirm our reported associations and further explore and explain the pathways through which they operate. However, the aforementioned studies used more descriptive statistical methods, assessed smaller parts of the spine and did not include assessments of inflammation. Our study applied statistical methods that disentangle the temporal sequence, control for inflammation, and handle the correlations between levels of data, and allow us to evaluate the link between two types of bony lesions at the VU level. We also explored the hypotheses in both directions and in multiple scenarios, looking not only to

1
2
3 associations on the same VU but also across VU levels. With this, our study has brought strong
4 arguments to a longstanding debate.
5

6
7 As mentioned above, all our models were adjusted for presence of inflammation. The effect of
8 vertebral body inflammation on the development of syndesmophytes was already known from
9 literature,[16-18] and was therefore added to our models. Although inflammation in the posterior
10 elements is well known in r-axSpA,[21, 22] its association with FJA was not previously reported.
11 Because of the known link between vertebral body inflammation and syndesmophytes, we
12 suspected inflammation in the posterior elements to have a similar effect on FJA and added it to the
13 models that used FJA as the outcome. In our study population, inflammation in the posterior
14 elements was most frequently present in the thoracic spine, with almost no presence in the cervical
15 and lower lumbar spine and little presence in the upper lumbar spine (Supplementary Figure S1).
16 Although the association between inflammation in the posterior elements and FJA was not the
17 primary aim of the study, statistically significant positive associations were found in almost half of
18 the models adjusted for it (Table 1 and 2). These results show that, in our multivariable models,
19 developing FJA is two times as likely when there is inflammation in the posterior elements two years
20 prior.
21
22

23
24 In conclusion, this is the first study to provide evidence pointing to a positive association between
25 FJA and syndesmophyte development and bridging syndesmophytes and FJA development, which
26 exists next to the associations between inflammation and bony lesions. Additionally, this is the first
27 study to report a positive association between inflammation in the posterior elements and FJA. With
28 these results we can imagine a pathway to pathological bone formation in r-axSpA in which
29 inflammation leads to bone formation, after which bone formation leads to additional bone
30 formation. Previous studies have reported several risk factors for syndesmophyte development but
31 have not taken into account other bone formation in the spine. We show that FJA is connected to
32 syndesmophyte development and should therefore also be considered and measured when studying
33 structural damage in the spine.
34
35

36 **Data availability statement**

37 The data underlying this article will be shared on reasonable request to the corresponding author.
38

39 **Funding**

40 This work was supported by Reuma Nederland and Novartis
41

42 **Conflicts of interest**

43 AS reports consulting fees from UCB, payment from Novartis, and support for attending meetings
44 from Lilly and UCB; DvdH reports financial support from the Dutch Rheumatism Association for the
45 SIAS study, consulting fees for Abbvie, Gilead, Glaxo-Smith-Kline, Lilly, Novartis and UCB, and a role
46 as director of Imaging Rheumatology bv.; FvG reports grants from Stichting vrienden van Sole Mio,
47 Stichting ASAS, Jacobus Stichting, Novartis and UCB and fees from Novartis and personal fees from
48 MSD, AbbVie, Bristol Myers Squibb, Eli-Lilly; MR reports grants for the ASAS CLASSIC study, outside
49 the submitted work; RL reports consulting fees and speaker honoraria from Abbvie, Amgen, BMS,
50 GSK, Jansen, Eli-Lilly, Novartis, Pfizer and UCB, participation on the data safety monitoring board for
51 a trial outside the submitted work and a role as chair of quality of care for EULAR; The other authors
52 have declared no conflicts of interest.
53
54
55
56
57
58
59
60

References

- 1 Navarro-Compán V, Sepriano A, El-Zorkany B, Van der Heijde D. Axial Spondyloarthritis. *Ann Rheum Dis* 2021;80(12):1511-21.
- 2 Braun J, van der Heijde D. Imaging and scoring in ankylosing spondylitis. *Best Pract Res Clin Rheumatol* 2002;16(4):573-604.
- 3 Reijnierse M. Radiographic/MR imaging correlation of paravertebral ossifications in ligaments and bony vertebral outgrowths. *Magn Reson Imaging Clin N Am* 2019;27(4):641-59.
- 4 Baraliakos X, Listing J, Rudwaleit M, Haibel H, Brandt J, Sieper J, et al. Progression of radiographic damage in patients with ankylosing spondylitis: defining the central role of syndesmophytes. *Ann Rheum Dis* 2007;66:910-5.
- 5 Maas F, Spoorenberg A, Brouwer E, van der Veer E, Bootsma H, Bos R, et al. Radiographic damage and progression of the cervical spine in ankylosing spondylitis patients treated with TNF- α inhibitors: Facet joints vs. vertebral bodies. *Semin Arthritis Rheum* 2017;46(5):562-8.
- 6 Stal R, van Gaalen F, Sepriano A, Braun J, Reijnierse M, van den Berg R, et al. Facet joint ankylosis in r-axSpA: detection and 2-year progression on whole spine low-dose CT and comparison with syndesmophyte progression. *Rheumatology (Oxford)* 2020;59(12):3776-83.
- 7 de Vlam K, Mielants H, Veys EM. Involvement of the zygapophyseal joint in ankylosing spondylitis: Relation to the bridging syndesmophyte. *J Rheumatol* 1999;26(8):1738-45.
- 8 Tan S, Yao J, Flynn JA, Yao L, Ward MM. Zygapophyseal Joint Fusion in Ankylosing Spondylitis Assessed by Computed Tomography: Associations with Syndesmophytes and Spinal Motion. *J Rheumatol* 2017;44(7):1004-10.
- 9 Braun J, Baraliakos X, Golder W, Hermann K, Listing J, Brandt J, et al. Analysing chronic spinal changes in ankylosing spondylitis: a systematic comparison of conventional x rays with magnetic resonance imaging using established and new scoring systems. *Ann Rheum Dis* 2004;63:1046-55.
- 10 Ramiro S, Claudepierre P, Sepriano A, van Lunteren M, Molto A, Feydy A, et al. Which scoring method depicts spinal radiographic damage in early axial spondyloarthritis best? Five-year results from the DESIR cohort. *Rheumatology (Oxford)* 2018;57:1991-2000.
- 11 Ramiro S, van Tubergen A, Stolwijk C, Landewé R, van de Bosch F, Dougados M, et al. Scoring radiographic progression in ankylosing spondylitis: should we use the modified Stoke Ankylosing Spondylitis Spine Score (mSASSS) or the Radiographic Ankylosing Spondylitis Spinal Score (RASSS)? *Arthritis Res Ther* 2013;15(1):R14.
- 12 Maas F, Arends S, Brouwer E, Bootsma H, Bos R, Wink FR, et al. Incorporating assessment of the cervical facet joints in the modified Stoke ankylosing spondylitis spine score is of additional value in the evaluation of spinal radiographic outcome in ankylosing spondylitis. *Arthritis Res Ther* 2017;19(1).
- 13 Tan S, Ward MM. Computed tomography in axial spondyloarthritis. *Curr Opin Rheumatol* 2018;30(4):334-9.
- 14 de Bruin F, de Koning A, van den Berg R, Baraliakos X, Braun J, Ramiro S, et al. Development of the CT Syndesmophyte Score (CTSS) in patients with ankylosing spondylitis: data from the SIAS cohort. *Ann Rheum Dis* 2018;77(3):371-7.
- 15 de Koning A, de Bruin F, van den Berg R, Ramiro S, Baraliakos X, Braun J, et al. Low-dose CT detects more progression of bone formation in comparison to conventional radiography in patients with ankylosing spondylitis: results from the SIAS cohort. *Ann Rheum Dis* 2018;77(2):293-9.
- 16 Tan S, Wang R, Ward MM. Syndesmophyte growth in ankylosing spondylitis. *Curr Opin Rheumatol* 2015;27(4):326-32.
- 17 Machado P, Baraliakos X, van der Heijde D, Braun J, Landewé R. MRI vertebral corner inflammation followed by fat deposition is the strongest contributor to the development of new bone at the same vertebral corner: a multilevel longitudinal analysis in patients with ankylosing spondylitis. *Ann Rheum Dis* 2016;75:1486-93.

1
2
3 18 van der Heijde D, Machado P, Braun J, Hermann K, Baraliakos X, Hsu B, et al. MRI inflammation at
4 the vertebral unit only marginally predicts new syndesmophyte formation: a multilevel analysis in
5 patients with ankylosing spondylitis. *Ann Rheum Dis* 2012;71(3):369-73.

6 19 Maksymowych WP, Inman RD, Salonen D, Dhillon SS, Krishnananthan R, Stone M, et al.
7 Spondyloarthritis Research Consortium of Canada magnetic resonance imaging index for assessment
8 of spinal inflammation in ankylosing spondylitis. *Arthritis Rheum* 2005;53(4):502-9.

9 20 Sepriano A, Ramiro S, van der Heijde D, Dougados M, Claudepierre P, Feydy A, et al. Integrated
10 longitudinal analysis does not compromise precision and reduces bias in the study of imaging
11 outcomes: A comparative 5-year analysis in the DESIR cohort. *Semin Arthritis Rheum* 2020;50:1394-
12 9.

13 21 Lee S, Lee J, Hwang J, Shin J, Kim T-H, Kim S-H. Clinical importance of inflammatory facet joints of
14 the spine in ankylosing spondylitis: a magnetic resonance imaging study. *Scand J Rheumatol*
15 2016;45(6):491-8.

16 22 Bochkova A, Levshakova AV, Bunchuk NV, Braun J. Spinal inflammation lesions as detected by
17 magnetic resonance imaging in patients with early ankylosing spondylitis are more often observed in
18 posterior structures of the spine. *Rheumatology (Oxford)* 2010;49(4):749-55.

VU	Segment	≥1 bridging syndesmophyte at BL		≥1 FJA at BL	
		Reader 1	Reader 2	Reader 1	Reader 2
1	Cervical	22%	27%	12%	34%
2		25%	25%	14%	21%
3		25%	29%	23%	20%
4		33%	34%	NA	NA
5		20%	29%	NA	NA
6		25%	23%	NA	NA
7	Thoracic	27%	31%	NA	NA
8		33%	39%	23%	28%
9		49%	47%	27%	33%
10		55%	59%	26%	37%
11		45%	55%	22%	35%
12		47%	47%	26%	29%
13		57%	57%	29%	29%
14		55%	57%	24%	31%
15		49%	55%	24%	33%
16		55%	59%	25%	37%
17		53%	49%	27%	29%
18		48%	48%	28%	36%
19	Lumbar	26%	28%	16%	24%
20		28%	24%	12%	22%
21		22%	22%	12%	14%
22		20%	22%	10%	20%
23		12%	12%	10%	12%

Figure 1: Occurrences of bridging syndesmophytes and facet joint ankylosis at baseline

Numbers represent the percentage of patients with the event (all coded binary), per vertebral unit, per reader. Maximum number of total patients per cell is 51 patients, numbers can be lower due to missing scores. Facet joint ankylosis scores at vertebral units 4-7 are excluded due to poor visibility. VU1 = C2-C3; VU23 = L5-S1. VU, vertebral unit; FJA, facet joint ankylosis; BL, baseline; NA, not applicable




Figure 2: Presence of bridging syndesmophytes and facet joint ankylosis at baseline on the patient level

The figure shows the presence of bridging syndesmophytes and facet joint ankylosis at baseline in each individual patient. The plot is ordered by increasing number of VUs with bridging syndesmophytes, and information on syndesmophytes and facet joint ankylosis is linked per patient (e.g. patient at the 0.4 probability of syndesmophytes has 5 bridging syndesmophytes and 0 ankylosed facet joints). Scores are the mean of both readers of the number of vertebral units with at least one bridging syndesmophyte and the number of vertebral units with at least one ankylosed facet joint. The range for both scores is 0-19. VU, vertebral unit.

Table 1: Multivariable models association between FJA and syndesmophyte development two years later and bridging syndesmophytes and the development of FJA two years later

	N=51 OR (95% CI)
Effect of FJA on increase syndesmophytes (hypothesis 1A)	
Baseline FJA	1.87 (1.20-2.92)
Baseline vertebral body inflammation	1.88 (1.28-2.76)
Effect of FJA on presence syndesmophytes (hypothesis 1B)	
Baseline FJA	1.69 (0.88-3.22)
Baseline vertebral body inflammation	2.32 (1.40-3.85)
Effect of bridging syndesmophytes on increase FJA (hypothesis 2A)	
Baseline bridging syndesmophyte	3.55 (2.03-6.21)
Baseline posterior elements inflammation	2.21 (0.73-6.65)
Effect of bridging syndesmophytes on presence FJA (hypothesis 2B)	
Baseline bridging syndesmophyte	3.30 (2.14-5.09)
Baseline posterior elements inflammation	2.49 (1.20-5.16)

Multivariate model results for the effects of facet joint ankylosis at baseline on the development of syndesmophytes 2 years later (hypothesis 1) and for the effects of bridging syndesmophytes at baseline on the development of facet joint ankylosis two years later (hypothesis 2). Two models are made per hypothesis. Model A looks at the effect of the predictor on the number of newly developed bony lesions (syndesmophyte or facet joint ankylosis) in a vertebral unit regardless of the presence of lesions at baseline. Model B looks at the development of a new lesion (syndesmophyte or facet joint ankylosis) on a vertebral unit adjusting for the presence of lesions at baseline. Vertebral units 4-7 are excluded due to poor visibility of the facet joints. Inflammation is defined as presence of an inflammatory lesion on the vertebral body of a vertebral unit by 2/3 MRI readers and as presence of an inflammatory lesion in any of the posterior elements of a vertebral unit (pedicles, processes, facet joints and soft tissues in vertebral units 7-23; only pedicles and soft tissues in vertebral units 1-6) by 2/3 MRI readers. FJA, facet joint ankylosis.

Table 2: Multivariable model associations for adjacent-VU analyses of FJA and syndesmophyte development two years later and bridging syndesmophytes and FJA development two years later

	1 VU-level shifted N=51 OR (95% CI)	2 VU-levels shifted N=51 OR (95% CI)
Effect of FJA on increase syndesmophytes on the VU above (hypothesis 1A)		
Baseline FJA	1.96 (1.23-3.14)	1.29 (0.79-2.10)
Baseline vertebral body inflammation	1.82 (1.24-2.67)	1.93 (1.30-2.86)
Effect of FJA on increase syndesmophytes on the VU below (hypothesis 1A)		
Baseline FJA	1.55 (0.99-2.43)	1.85 (1.25-2.74)
Baseline vertebral body inflammation	1.80 (1.19-2.71)	1.78 (1.20-2.63)
Effect of FJA on presence syndesmophytes on the VU above (hypothesis 1B)		
Baseline FJA	2.01 (1.06-3.82)	1.19 (0.53-2.71)
Baseline vertebral body inflammation	2.39 (1.45-3.95)	2.40 (1.42-4.05)
Effect of FJA on presence syndesmophytes on the VU below (hypothesis 1B)		
Baseline FJA	1.89 (1.03-3.45)	2.19 (1.12-4.31)
Baseline vertebral body inflammation	2.61 (1.53-4.44)	2.16 (1.26-3.68)
Effect bridging syndesmophyte on increase FJA on the VU above (hypothesis 2A)		
Baseline bridging syndesmophyte	2.56 (1.23-5.31)	1.70 (0.84-3.45)
Baseline posterior elements inflammation	1.97 (0.64-6.06)	2.05 (0.73-5.75)
Effect bridging syndesmophyte on increase FJA on the VU below (hypothesis 2A)		
Baseline bridging syndesmophyte	3.11 (1.88-5.12)	2.98 (1.97-4.50)
Baseline posterior elements inflammation	2.21 (0.75-6.54)	1.98 (0.68-5.79)
Effect bridging syndesmophyte on presence FJA on the VU above (hypothesis 2B)		
Baseline bridging syndesmophyte	1.81 (1.01-3.25)	1.47 (0.93-2.32)
Baseline posterior elements inflammation	2.30 (1.12-4.71)	2.29 (1.19-4.40)
Effect bridging syndesmophyte on presence FJA on the VU below (hypothesis 2B)		
Baseline bridging syndesmophyte	3.04 (1.94-4.77)	3.13 (1.90-5.16)
Baseline posterior elements inflammation	2.45 (1.20-5.00)	2.51 (1.22-5.16)

Multivariable model results for the effects of facet joint ankylosis at baseline on the development of syndesmophytes 2 years later on an adjacent vertebral unit (hypothesis 1) and for the effects of bridging syndesmophytes at baseline on the development of facet joint ankylosis two years later on an adjacent vertebral unit (hypothesis 2). Two types of models are made per hypothesis. Model A looks at the effect of the predictor on the number of newly developed bony lesions (syndesmophyte or facet joint ankylosis) in an adjacent vertebral unit regardless of the presence of lesions at baseline

Model B looks at the development of a new lesion (syndesmophyte or facet joint ankylosis) in an adjacent vertebral unit adjusting for the presence of lesions at baseline. The shifting of vertebral units is introduced only between the predictor and the outcome variable, i.e. inflammation at baseline is still added as a predictor for syndesmophytes at follow-up on the same vertebral unit. Facet joint ankylosis scores for vertebral units 4-7 are excluded due to poor visibility of the facet joints. Inflammation is defined as presence of an inflammatory lesion on the vertebral body of a vertebral unit by 2/3 MRI readers and as presence of an inflammatory lesion in any of the posterior elements of a vertebral unit (pedicles, processes, facet joints and soft tissues in vertebral units 7-23; only pedicles and soft tissues in vertebral units 1-6) by 2/3 MRI readers. FJA, facet joint ankylosis. VU, vertebral unit; FJA, facet joint ankylosis; BL, baseline.

Table 3: multivariable models associations of dose-response effects between FJA and syndesmophytes

	N=51 OR (95% CI)
Effect of FJA on increase syndesmophytes (hypothesis 1A)	
Baseline FJA – no ankylosis	Ref.
Baseline FJA – 1 ankylosed facet joint	1.32 (0.54-3.22)
Baseline FJA – 2 ankylosed facet joints	2.15 (1.24-3.27)
Baseline vertebral body inflammation	1.87 (1.27-2.75)
Effect of FJA on presence syndesmophytes (hypothesis 1B)	
Baseline FJA – no ankylosis	Ref.
Baseline FJA – 1 ankylosed facet joint	1.17 (0.25-5.35)
Baseline FJA – 2 ankylosed facet joints	1.86 (0.94-3.71)
Baseline vertebral body inflammation	2.30 (1.40-3.79)
Effect of bridging syndesmophytes on increase FJA (hypothesis 2A)	
Baseline bridging syndesmophyte – 0 affected quadrants	Ref.
Baseline bridging syndesmophyte – 1 affected quadrant	2.22 (0.90-5.46)
Baseline bridging syndesmophyte – 2 affected quadrants	3.09 (1.30-7.36)
Baseline bridging syndesmophyte – 3 affected quadrants	4.83 (2.03-11.52)
Baseline bridging syndesmophyte – 4 affected quadrants	6.43 (2.95-14.03)
Baseline posterior elements inflammation	2.14 (0.69-6.61)
Effect of bridging syndesmophytes on presence FJA (hypothesis 2B)	
Baseline bridging syndesmophyte – 0 affected quadrants	Ref.
Baseline bridging syndesmophyte – 1 affected quadrant	2.35 (1.27-4.34)
Baseline bridging syndesmophyte – 2 affected quadrants	3.29 (1.60-6.77)
Baseline bridging syndesmophyte – 3 affected quadrants	3.23 (1.23-8.45)
Baseline bridging syndesmophyte – 4 affected quadrants	5.54 (3.14-9.80)
Baseline posterior elements inflammation	2.51 (1.22-5.15)

Multivariate model results for the effects of facet joint ankylosis at baseline on the development of syndesmophytes 2 years later (hypothesis 1) and for the effects of bridging syndesmophytes at baseline on the development of facet joint ankylosis two years later (hypothesis 2). Predictors are categorized in the extend of the bony lesion present: FJA in 0-2 joints and bridging syndesmophytes in 0-4 quadrants. In all models, the category with no presence of the predictor (i.e. zero ankylosed

1
2
3 facet joints or 0 affected quadrants) is the reference for the other categories. Two models are made
4 per hypothesis. Model A looks at the effect of the predictor on the number of newly developed bony
5 lesions (syndesmophyte or facet joint ankylosis) in a vertebral unit regardless of the presence of
6 lesions at baseline. Model B looks at the development of a new lesion (syndesmophyte or facet joint
7 ankylosis) on a vertebral unit adjusting for the presence of lesions at baseline. Vertebral units 4-7 are
8 excluded due to poor visibility of the facet joints. Inflammation is defined as presence of an
9 inflammatory lesion on the vertebral body of a vertebral unit by 2/3 MRI readers and as presence of
10 an inflammatory lesion in any of the posterior elements of a vertebral unit (pedicles, processes, facet
11 joints and soft tissues in vertebral units 7-23; only pedicles and soft tissues in vertebral units 1-6) by
12 2/3 MRI readers. FJA, facet joint ankylosis.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60